

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 209/08, 209/30, 209/40, A61K 31/404, A61P 5/36		A1	(11) International Publication Number: WO 00/66555
			(43) International Publication Date: 9 November 2000 (09.11.00)
(21) International Application Number: PCT/US00/11630		<p>Lin [CN/US]; 7794 Roan Road, San Diego, CA 92129 (US). JONES, Todd, K. [US/US]; 546 Marview Drive, Solana Beach, CA 92075 (US). MARSCHKE, Keith, B. [US/US]; 12565 Escala Drive, San Diego, CA 92128 (US). TEGLEY, Christopher, M. [US/US]; 478 Thunderhead Street, Thousand Oaks, CA 91360 (US).</p> <p>(74) Agents: KODROFF, Cathy, A. et al.; Howson and Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).</p> <p>(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
(22) International Filing Date: 1 May 2000 (01.05.00)			
(30) Priority Data: 60/172,259 4 May 1999 (04.05.99) US 09/552,033 19 April 2000 (19.04.00) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/552,033 (CON) Filed on 19 April 2000 (19.04.00)			
(71) Applicants (for all designated States except US): AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07840 (US). LIGAND PHARMACEUTICALS, INC. [US/US]; 10275 Science Center Drive, San Diego, CA 92121 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): FENSOME, Andrew [GB/US]; 779 Trephanny Lane, Wayne, PA 19087 (US). ZHANG, Puwen [CN/US]; 7004 Oriole Court, Audubon, PA 19403 (US). KOKO, Marci, C. [US/US]; 220 West Langhorne Avenue, Bethlehem, PA 19087 (US). ZHI,			
(54) Title: THIO-OXINDOLE DERIVATIVES			
<div style="display: flex; justify-content: space-around; align-items: center;"><div style="text-align: center;"><p>(1)</p></div><div style="text-align: center;"><p>(2)</p></div></div>			
(57) Abstract			
<p>This invention relates to compounds which are agonists of the progesterone receptor which have general structures (1) or (2) wherein the substituents are as defined; or a pharmaceutically acceptable salt thereof, as well as methods of using these compounds to induce contraception or treat progesterone-related carcinomas and adenocarcinomas.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

THIO-OXINDOLE DERIVATIVES

Field of the Invention

5 This invention relates to compounds which are agonists of the progesterone receptor, their preparation and utility.

Background of the Invention

 Intracellular receptors (IR) form a class of structurally related gene regulators
10 known as "ligand dependent transcription factors" (R. M. Evans, *Science*, **240**, 889, 1988). The steroid receptor family is a subset of the IR family, including progesterone receptor (PR), estrogen receptor (ER), androgen receptor (AR), glucocorticoid receptor (GR), and mineralocorticoid receptor (MR).

 The natural hormone, or ligand, for the PR is the steroid progesterone, but
15 synthetic compounds, such as medroxyprogesterone acetate or levonorgestrel, have been made which also serve as ligands. Once a ligand is present in the fluid surrounding a cell, it passes through the membrane *via* passive diffusion, and binds to the IR to create a receptor/ligand complex. This complex binds to specific gene promoters present in the cell's DNA. Once bound to the DNA the complex modulates
20 the production of mRNA and protein encoded by that gene.


 A compound that binds to an IR and mimics the action of the natural hormone is termed an agonist, whilst a compound which inhibits the effect of the hormone is an antagonist.

 PR agonists (natural and synthetic) are known to play an important role in the
25 health of women. PR agonists are used in birth control formulations, typically in the presence of an ER agonist, alternatively they may be used in conjunction with a PR antagonist. ER agonists are used to treat the symptoms of menopause, but have been associated with a proliferative effect on the uterus which can lead to an increased risk of uterine cancers. Co-administration of a PR agonist reduces/ablates that risk.


- 2 -

The compounds of this invention have been shown to act as competitive inhibitors of progesterone binding to the PR and act as agonists. These compounds may be used for contraception and post menopausal hormone replacement therapy.

Jones, *et al*, described in U.S. Patent No. 5,688,810 the PR antagonist dihydroquinoline A.


**A**

Jones, *et al*, described the enol ether B (U.S. Patent No. 5,693,646) as a PR ligand.


**B**

Jones, *et al*, described compound C (U.S. Patent No. 5,696,127) as a PR ligand.

- 3 -




Zhi, *et al*, described lactones **D**, **E** and **F** as PR antagonists (J. Med. Chem., 41, 291, 1998).




Zhi, *et al*, described the ether **G** as a PR antagonist (J. Med. Chem., 41, 291, 1998).

- 4 -

**G**


Combs, *et al.*, disclosed the amide **H** as a ligand for the PR (*J. Med. Chem.*,
38, 4880, 1995).

5


**H**

Perlman, *et al.*, described the vitamin D analog **I** as a PR ligand (*Tet. Letters*,
10 35, 2295, 1994).


- 5 -

**I**

Hamann, *et al*, described the PR antagonist **J** (*Ann. N.Y. Acad. Sci.*, **761**, 383, 1995).


**J**

Chen, *et al*, described the PR antagonist **K** (Chen, *et al*, POI-37, 16th Int. Cong. Het. Chem., Montana, 1997).

**K**


- 6 -

Kurihari, *et al.*, described the PR ligand **L** (*J. Antibiotics*, **50**, 360, 1997).

**L**


5

Kuhla, et al, taught the oxindole **M** as a cardiotonic (WO 86/03749).

**M**

10


Weber, described the oxindole **N** for cardiovascular indications (WO 91/06545).

**N**

15


Fischer, et al, claim a preparation for making compounds which include the generic structure **O** (U.S. Patent No. 5,453,516).

- 7 -




R = various

- 5 Singh, et al, described the PDE III inhibitor P (*J. Med. Chem.*, 37, 248, 1994).




Andreani, et al, described the cytotoxic agent Q (*Acta. Pharn. Nord.*, 2, 407, 1990).

10




- Binder, et al, described structure R which is an intermediate for preparing COX II
15 inhibitors (WO 97/13767).

- 8 -



Walsh (A.H. Robins) described the oxindole S as an intermediate (U.S. Patent No.
5 4,440,785, U.S. Patent No. 4,670,566).




R1 = F, Cl, Br, alkyl, NH₂

R2 = alkyl, alkoxy, F, Cl, NH₂, CF₃

10


Bohm, et al, claim the oxindole T as cardiovascular agents (WO 91/06545).



15


Bohm, et al, include the generic structure U (WO 91/04974).

- 9 -




JP 63112584 A contains the generic structure V:

5




Boar, et al, described the dioxolane W as an intermediate for preparation of acetylcholinesterase inhibitors (WO 93/12085 A1).

10



- 10 -


Kende, et al, described methodology for preparing 3,3-substituted oxindoles, e.g. X, that was utilized in the present invention (Synth. Commun., 12, 1, 1982).



X

Description of the Invention

This invention provides compounds of the formulae 1 or 2:



wherein:

R_1 and R_2 are chosen independently from the group of H, alkyl, substituted alkyl; OH; O(alkyl); O(substituted alkyl); OAc; aryl; optionally substituted aryl; heteroaryl; optionally substituted heteroaryl; alkylaryl; alkylheteroaryl; 1-propynyl; or 3-propynyl:

or R_1 and R_2 are joined to form a ring comprising one of the following:

$-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$; $-\text{CH}_2\text{CH}_2\text{CMe}_2\text{CH}_2\text{CH}_2-$; $-\text{O}(\text{CH}_2)_m\text{CH}_2-$; $\text{O}(\text{CH}_2)_p\text{O}-$;
 $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$; or $-\text{CH}_2\text{CH}_2\text{N}(\text{H or alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

- 11 -

p is an integer from 1 to 4;

or R₁ and R₂ together comprise a double bond to one of the following:

CMe₂; C(cycloalkyl), O, C(cyloether).


R₃ is selected from H, OH, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₆ alkenyl, alkynyl or substituted alkynyl, or COR^A;

R^A is selected from H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R₄ is selected from H, halogen, CN, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₁ to C₆ alkoxy, substituted C₁ to C₆ alkoxy, C₁ to C₆ aminoalkyl, or substituted C₁ to C₆ aminoalkyl;

R^S is selected from the groups a), b) or c):

a) R^S is a trisubstituted benzene ring containing the substituents X, Y and Z as shown below:



15

X is selected from halogen, OH, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkyl, substituted C₁ to C₃ thioalkyl, S(O)alkyl, S(O)₂alkyl, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 or 6 membered heterocyclic ring containing 1 to 3 heteroatoms, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, COR^B, OCOR^B, NR^CCOR^B;

R^B is selected from H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^C is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

Y and Z are independently selected from H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, or C₁ to C₃ thioalkyl;

25

- 12 -

or

b) R^5 is a five or six membered heterocyclic ring with 1, 2, or 3 heteroatoms selected from O, S, SO, SO_2 or NR^6 and containing one or two independent substituents from the group of H, halogen, CN, NO_2 and C_1 to C_3 alkyl, C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, COR^D , or $NR^E COR^D$;

R^D is H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, or substituted C_1 to C_3 aminoalkyl;

R^E is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

R^6 is H, or C_1 to C_3 alkyl; or

c) R^5 is an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, the moiety being optionally substituted by from 1 to 3 substituents selected from halogen, lower alkyl, CN, NO_2 , lower alkoxy, or CF_3 ;


Q^1 is S, NR_7 , CR_8R_9 ;

R_7 is selected from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, acyl, substituted acyl, aroyl, substituted aroyl, SO_2CF_3 , OR^{11} or $NR^{11}R^{12}$;

R_8 and R_9 are independent substituents selected from the group of H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN, or CO_2R_{10} ,


R_{10} is C_1 to C_3 alkyl; or

CR_8R_9 comprises a six membered ring as shown by the structure below



Q^2 is selected from the moieties:

- 13 -



R^{11} , R^{12} and R^{13} are independently selected from H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, aroyl or substituted aroyl or sulfonyl;

5

or a pharmaceutically acceptable salt thereof.


A preferred list of substituents represented by R^{11} , R^{12} and R^{13} in groups of the compounds described herein are H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, $-\text{C}(\text{O})-$ (C_1 to C_6 alkyl), $-\text{S}(\text{O})_2-(\text{C}_1$ to C_6 alkyl), phenyl or benzyl.

10


It will be understood that this invention includes all tautomeric forms of the compounds, chemical formulae and substituents described herein.

Two preferred sets of compounds of this invention is depicted by structures 2 and 3, respectively:

15



each wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below




- 14 -

X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

Another preferred group of formula 2 are those wherein R₅ is a five membered ring with the structure shown below




wherein:

- 10 U is O, S, or NR₆;
 R₆ is H, or C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;
 X' is selected from halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkyl, or C₁ to C₃ alkoxy;
 Y' is from the group of H, F or C₁ to C₄ alkyl;
 15 or a pharmaceutically acceptable salt thereof.

A further preferred subgroup of the compounds above are those in which R₅ is a thiophene or furan ring substituted by X' and Y', as described above.

A further preferred subgroup group of compounds of formulas 2 and 3 are those wherein R₅ is a six membered ring with the structure:



20

wherein X' is N or CX²,

- 15 -

X^2 is halogen, CN, CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CONalkyl_2 , CSNalkyl_2 or NO_2 ;


Q^1 is S, NR_7 , CR_8R_9 ;

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl,
 5 C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl,
 heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6
 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl,
 aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN , CO_2R_{10} ,


10 R_{10} is C_1 to C_3 alkyl;

CR_8R_9 are within a six membered ring as shown by the structure below




or pharmaceutically acceptable salt thereof.

15 Still another preferred group of these compounds includes those having the
 general formulae:



each wherein R_5 is a disubstituted benzene ring containing the substituents X
 and Y as shown below


- 16 -




X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy;

5 Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

A further preferred subgroup group of compounds of formulae:



10 are those wherein R₅ is a six membered ring with the structure:



wherein X¹ is N or CX²,

X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂ or NO₂;

15 Q² is as defined above;


R₇ is from the group including CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO₂CF₃;

- 17 -

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN , CO_2R_{10} ,

R_{10} is C_1 to C_3 alkyl;



5 CR_8R_9 are within a six membered ring as shown by the structure below



or pharmaceutically acceptable salt thereof.


A further preferred set of compounds of this invention is depicted by structure

10 4,



Wherein R_{14} is chosen from the group H, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, substituted sulfonyl.

15 Wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below




- 18 -

X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNHOH, CNH₂NOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy;

- 5 Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

Another preferred group of formula 4 are those wherein R₅ is a five membered ring with the structure shown below



- 10 wherein:

U is O, S, or NR₆;

R₆ is H, or C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from halogen, CN, NO₂, CONH₂, CNHNHOH, CNH₂NOH, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkyl, or C₁ to C₃ alkoxy;


- 15

Y' is from the group of H, F or C₁ to C₄ alkyl;

or a pharmaceutically acceptable salt thereof.

A further preferred subgroup of the compounds above are those in which R₅ is a thiophene or furan ring substituted by X' and Y', as described above.

- 20 A further preferred subgroup group of compounds of formula 4 are those wherein R₅ is a six membered ring with the structure:



- 19 -

wherein X^1 is N or CX^2 ,

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl_2$, $CSNalkyl_2$ or NO_2 ;


R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl,
 5 C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl,
 heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6
 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl,
 aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN CO_2R_{10} ,


10 R_{10} is C_1 to C_3 alkyl;
 or pharmaceutically acceptable salt thereof.

A further preferred set of compounds of this invention is depicted by structure
 5,

15



Wherein R_5 is a disubstituted benzene ring containing the substituents X and
 Y as shown below



20 X is selected from halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$,
 $CONalkyl_2$, $CSNalkyl_2$, $CNHNOH$, C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3


- 20 -

perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C_1 to C_3 thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, or C_1 to C_3 thioalkyl;

5 or a pharmaceutically acceptable salt thereof.

Another preferred group of formula 5 are those wherein R_5 is a five membered ring with the structure shown below



wherein:

10 U is O, S, or NR_6 ;

R_6 is H, or C_1 to C_3 alkyl, or C_1 to C_4 CO_2 alkyl;


X' is selected from halogen, CN, NO_2 , $CONH_2$, $CSNH_2$, $CONH$ alkyl, $CSNH$ alkyl, CON alkyl₂, CSN alkyl₂, C_1 to C_3 alkyl, or C_1 to C_3 alkoxy;

Y' is from the group of H, F or C_1 to C_4 alkyl;

15 or a pharmaceutically acceptable salt thereof.

A further preferred subgroup of the compounds above are those in which R_5 is a thiophene or furan ring substituted by X' and Y' , as described above.

A further preferred subgroup group of compounds of formula 5 are those wherein R_5 is a six membered ring with the structure:



20

wherein X^1 is N or CX^2 ,

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONH$ alkyl, $CSNH$ alkyl, CON alkyl₂, CSN alkyl₂ or NO_2 ;


- 21 -

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN, CO_2R_{10} ,


R_{10} is C_1 to C_3 alkyl;
or pharmaceutically acceptable salt thereof.

A further preferred set of compounds of this invention is depicted by structure 6,



Wherein R_{15} is selected from the group H, Me, CO_2R , acyl, substituted acyl, aroyl, substituted aroyl, alkyl, substituted alkyl, CN.

Wherein R_6 is a disubstituted benzene ring containing the substituents X and Y as shown below




X is selected from halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl_2$, $CSNalkyl_2$, $CNHNOH$, C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C_1 to C_3 thioalkoxy;

- 22 -

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

- 5 Another preferred group of formula 6 are those wherein R₅ is a five membered ring with the structure shown below



wherein:

U is O, S, or NR₆;

R₆ is H, or C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;


- 10 X' is selected from halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkyl, or C₁ to C₃ alkoxy;

Y' is from the group of H, F or C₁ to C₄ alkyl;

or a pharmaceutically acceptable salt thereof.

- A further preferred subgroup of the compounds above are those in which R₅ is a thiophene or furan ring substituted by X' and Y', as described above.
- 15

A further preferred subgroup of compounds of formula 6 are those wherein R₅ is a six membered ring with the structure:



wherein X¹ is N or CX²,

- 20 X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂ or NO₂;


- 23 -

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;


R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN CO_2R_{10} ,

R_{10} is C_1 to C_3 alkyl;
or pharmaceutically acceptable salt thereof.

A further preferred set of compounds of this invention is depicted by structure 7,




Wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below



X is selected from halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl$, $CSNalkyl$, $CNHNOH$, C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C_1 to C_3 thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, or C_1 to C_3 thioalkyl;
or a pharmaceutically acceptable salt thereof.

Another preferred group of formula 7 are those wherein R_5 is a five membered ring with the structure shown below



wherein:

U is O, S, or NR_6 ;

5 R_6 is H, or C_1 to C_3 alkyl, or C_1 to C_4 CO_2 alkyl;


X' is selected from halogen, CN, NO_2 , $CONH_2$, $CSNH_2$, $CONH$ alkyl, $CSNH$ alkyl, CON alkyl₂, CSN alkyl₂, C_1 to C_3 alkyl, or C_1 to C_3 alkoxy;

Y' is from the group of H, F or C_1 to C_4 alkyl;

or a pharmaceutically acceptable salt thereof.

10 A further preferred subgroup of the compounds above are those in which R_5 is a thiophene or furan ring substituted by X' and Y' , as described above.

A further preferred subgroup group of compounds of formula 7 are those wherein R_5 is a six membered ring with the structure:



15 wherein X' is N or CX^2 ,

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONH$ alkyl, $CSNH$ alkyl, CON alkyl₂, CSN alkyl₂ or NO_2 ;

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl,

20 heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN CO_2R_{10} ,

R₁₀ is C₁ to C₃ alkyl;
or pharmaceutically acceptable salt thereof.

5 The compounds of this invention may contain an asymmetric carbon atom and some of the compounds of this invention may contain one or more asymmetric centers and may thus give rise to optical isomers and diastereomers. While shown without respect to stereochemistry in Formula 1 and 2 the present invention includes such optical isomers and diastereomers; as well as the racemic and resolved,
10 enantiomerically pure R and S stereoisomers; as well as other mixtures of the R and S stereoisomers and pharmaceutically acceptable salts thereof.

 The term "alkyl" is used herein to refer to both straight- and branched-chain saturated aliphatic hydrocarbon groups having 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms; "alkenyl" is intended to include both straight- and branched-chain
15 alkyl group with 1 or 2 carbon-carbon double bonds and containing 2 to 8 carbon atoms, preferably 2 to 6 carbon atoms; "alkynyl" group is intended to cover both straight- and branched-chain alkyl group with at least 1 or 2 carbon-carbon triple bonds and containing 2 to 8 carbon atoms, preferably 2 to 6 carbon atoms.

 The term "acyl" refers to a carbonyl substituent, including both straight- and
20 branched-chain saturated aliphatic hydrocarbon groups having 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms. The term "substituted acyl" refers to an acyl group as just described optionally substituted with from 1 to 6 groups chosen from the list halogen, CN, OH, and NO₂.

 The term "aroyl" also refers to a carbonyl substituent carrying a phenyl group
25 or a heteroaromatic group. The heteroaromatic groups of this include 2-, 3- or 4-pyridinyl, 2- and 3-furanyl, 2- or 3-thiophenyl, or 2- or 4-pyrimidinal. The term "substituted aroyl" also refers to an aroyl group as just described optionally substituted with from 1 to 6 groups chosen from the list halogen, CN, OH, and NO₂.

 The terms "substituted alkyl", "substituted alkenyl", and "substituted
30 alkynyl" refer to alkyl, alkenyl, and alkynyl as just described having one or more substituents from the group including halogen, CN, OH, NO₂, amino, aryl,

heterocyclic, substituted aryl, substituted heterocyclic, alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl, alkylcarboxy, alkylamino, arylthio. These substituents may be attached to any carbon of alkyl, alkenyl, or alkynyl group provided that the attachment constitutes a stable chemical moiety.

5 The term "aryl" is used herein to refer to an aromatic system which may be a single ring or multiple aromatic rings fused or linked together as such that at least one part of the fused or linked rings forms the conjugated aromatic system. The aryl groups include, but are not limited to, phenyl, naphthyl, biphenyl, anthryl, tetrahydronaphthyl, phenanthryl.

10 The term "substituted aryl" refers to aryl as just defined having 1 to 4 substituents from the group including halogen, CN, OH, NO₂, amino, alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl, alkylcarboxy, alkylamino, or arylthio.


 The term "heterocyclic" is used herein to describe a stable 4- to 7-membered
15 monocyclic or a stable multicyclic heterocyclic ring which is saturated, partially unsaturated, or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group including N, O, and S atoms. The N and S atoms may be oxidized. The heterocyclic ring also includes any multicyclic ring in which any of above defined heterocyclic rings is fused to an aryl ring. The
20 heterocyclic ring may be attached at any heteroatom or carbon atom provided the resultant structure is chemically stable. Such heterocyclic groups include, for example, tetrahydrofuran, piperidinyl, piperazinyl, 2-oxopiperidinyl, azepinyl, pyrrolidinyl, imidazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, isoxazolyl, morpholinyl, indolyl, quinolinyl, thienyl, furyl, benzofuranyl,
25 benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide, and isoquinolinyl.

 The term "substituted heterocyclic" is used herein to describe the heterocyclic just defined having 1 to 4 substituents selected from the group which includes halogen, CN, OH, NO₂, amino, alkyl, substituted alkyl, cycloalkyl, alkenyl, substituted alkenyl, alkynyl, alkoxy, aryloxy, substituted alkyloxy, alkylcarbonyl,
30 alkylcarboxy, alkylamino, or arylthio.

- 27 -

- The term "thioalkyl" is used herein to refer to the SR group, where R is alkyl or substituted alkyl, containing 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms. The term "alkoxy" is used herein to refer to the OR group, where R is alkyl or substituted alkyl, containing 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms. The term "aryloxy" is used herein to refer to the OR group, where R is aryl or substituted aryl, as defined above. The term "alkylcarbonyl" is used herein to refer to the RCO group, where R is alkyl or substituted alkyl, containing 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms. The term "alkylcarboxy" is used herein to refer to the COOR group, where R is alkyl or substituted alkyl, containing 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms. The term "aminoalkyl" refers to both secondary and tertiary amines wherein the alkyl or substituted alkyl groups, containing 1 to 8 carbon atoms, preferably 1 to 6 carbon atoms, which may be either the same or different and the point of attachment is on the nitrogen atom. The term "halogen" refers to Cl, Br, F, or I.
- The compounds of this invention may be prepared according to the methods described below.

Scheme 1




According to scheme 1, commercially available oxindole 3 is treated with a strong organo-metallic base (e.g. butyl lithium, lithium diisopropylamide, potassium hexamethyldisilazide) in an inert solvent (e.g. THF, diethyl ether) under nitrogen at reduced temperature (ca. -20 °C) (Kende, et al, Synth. Commun., 12, 1, 1982) in the presence of lithium chloride or N,N,N',N'-tetramethylethylene-diamine. The resulting di-anion is then treated with excess electrophile such as an alkyl halide, preferably an iodide. If R₁ and R₂ are to be joined such as the product 4 contains a spirocycle at position 3, then the electrophile should be bifunctional, i.e. a diiodide. Subsequent bromination of 4 proceeds smoothly with bromine in acetic acid (an organic co-solvent such as dichloromethane may be added as required) in the presence of sodium acetate, to afford the aryl bromide 5. The bromide 5 is reacted with a palladium salt (e.g. tetrakis(triphenylphosphine)palladium(0) or palladium acetate), in a suitable solvent (e.g. THF, dimethoxyethane, acetone, ethanol or toluene) at room temperature under an inert atmosphere (argon, nitrogen). The mixture is then treated with an aryl or heteroaryl boronic acid or boronic acid ester and a base (sodium carbonate, triethylamine, potassium phosphate) in water or fluoride source (cesium fluoride) under anhydrous conditions. The required product 6 is then isolated and purified by standard means.

Reaction of the indoline-2-one derivative 6 with either Lawesson's reagent or phosphorous pentasulfide in a suitable organic solvent (pyridine, THF, dioxane, dimethoxyethane, dichloromethane, benzene, toluene, xylene) at a temperature between room temperature and the reflux temperature of the solvent provides access to the thiocarbonyl derivative 7. An additive such as sodium hydrogen carbonate may also be useful.

If R₁ and R₂ are different then the intermediate 4 is prepared by reacting the dianion of 3 with one equivalent of the electrophile R₁-X (X = leaving group e.g. iodine). The resultant mono-alkylated compound may then be isolated and re-subjected to the reaction conditions using R₂-X, or alternatively used *in-situ* for the second alkylation with R₂-X. Alternatively if the desired product 7 is to contain R₂ =

H, then the isolated mono-alkylated intermediate is taken through the subsequent steps.

Scheme 2




5 Other methodologies are also available for coupling the pendant aryl or heteroaryl group, Ar, to the oxindole platform, for example reaction of compound 5 with an aryl or heteroaryl stannane, aryl or heteroaryl zinc, or aryl or heteroaryl magnesium halide in the presence of a palladium or nickel catalyst (scheme 2). The required aryl or heteroaryl-metallic species described above are formed through
10 standard techniques.

Other functionalities can also be installed into the 3-position of the indoline platform according to scheme 3. Oxidation of the unsubstituted indoline 8, preferably under neutral or acidic conditions (e.g. selenium dioxide in dry dioxane at reflux) affords the isatin 9. Compound 9 may be further functionalized to provide a ketal 11
15 by treatment with an alcohol and acid catalyst under dehydrating conditions. Alternatively reaction of 9 with a second ketone under suitable conditions (piperidine in toluene at reflux; or TiCl₄/Zn in THF at reflux) affords alkylidene derivatives 11. Reaction of the isatin 9 with a grignard reagent or organolithium affords tertiary alcohols 12 (R = H). These alcohols may then be further functionalized by alkylation
20 or acylation procedures.

- 30 -


Scheme 3



Reaction of the indoline-2-one derivative 6 with either Lawesson's reagent or phosphorous pentasulfide in a suitable organic solvent (pyridine, THF, dioxane, dimethoxyethane, dichloromethane, benzene, toluene, xylene) at a temperature
5 between room temperature and the reflux temperature of the solvent provides access to the thiocarbonyl derivative 7. An additive such as sodium hydrogen carbonate may also be useful.

- 31 -


Scheme 4



- An alternative mode of preparation is to react compound 5 with either Lawesson's reagent or phosphorous pentasulfide in a suitable organic solvent (pyridine, THF, dioxane, dimethoxyethane, dichloromethane, benzene, toluene, xylene) at a temperature between room temperature and the reflux temperature of the solvent, under an inert atmosphere (nitrogen or argon) providing access to the thiocarbonyl derivative 13. Then reaction of bromide 13 in an anhydrous solvent (e.g. THF, Et₂O) with a strong base (sodium hydride preferred, sodium hexamethyldisilazide, potassium hydride) followed by reaction at reduced temperature (-50 to -20 °C) with n-butyllithium and *N,N,N,N'*-tetramethylethylenediamine followed after a suitable period of time by a trialkylborate (trimethyl or triisopropylborate) gives after acidic work-up the boronic acid 14 (scheme 4). Compound 14 may then be reacted under palladium catalyzed conditions (tetrakis(triphenylphosphine)palladium(0) or palladium acetate, base (NaHCO₃, Na₂CO₃, K₂CO₃, triethylamine, CsF) solvent (toluene/EtOH/water, THF/water, dimethoxyethane/water, anhydrous dimethoxyethane) with an aryl or heteroaryl bromide, aryl or heteroaryl iodide, aryl or heteroaryl trifluoromethane sulfonate or aryl or heteroaryl fluorosulfonate, to provide the desired compounds 7.

Alternatively reaction of compound **13** under palladium catalyzed conditions (tetrakis(triphenylphosphine)palladium(0) or palladium acetate, base (NaHCO_3 , Na_2CO_3 , K_2CO_3 , triethylamine, CsF) solvent (acetone/water, toluene/EtOH/water, THF/water, dimethoxyethane/water, anhydrous dimethoxyethane) with an aryl or
 5 heteroaryl bromide, aryl or heteroaryl iodide, aryl or heteroaryl trifluoromethane sulfonate or aryl or heteroaryl fluorosulfonate, to provide the desired compounds **7**.

Scheme 5



10 Treatment of the bromide **5** in an anhydrous solvent (e.g. THF, Et_2O) with a strong base (sodium hydride preferred, sodium hexamethyldisilazide, potassium hydride) followed by reaction at reduced temperature (-50 to -20 $^\circ\text{C}$) with *n*-butyllithium and *N,N,N,N'*-tetramethylethylenediamine followed after a suitable
 15 work-up the boronic acid **15** (scheme 5). Compound **15** may then be reacted under palladium catalyzed conditions (tetrakis(triphenylphosphine)palladium(0), base (NaHCO_3 , Na_2CO_3 , K_2CO_3 , triethylamine, CsF) solvent (toluene/EtOH/water, THF/water, dimethoxyethane/water, anhydrous dimethoxyethane) with an aryl or

- 33 -


heteroaryl bromide, aryl or heteroaryl iodide, aryl or heteroaryl trifluoromethane sulfonate or aryl or heteroaryl fluorosulfonate, to provide the desired compounds 6.

- 5 An alternative strategy would be to prepare an organo zinc or magnesium reagent from compound 5 and react it *in-situ* with an aryl or heteroaryl bromide, aryl or heteroaryl iodide, aryl or heteroaryl trifluoromethane sulfonate or aryl or heteroaryl fluorosulfonate, under palladium catalyzed conditions to afford compound 6. Such an organo zinc or magnesium species could be prepared by treatment of the bromide 57 in an anhydrous solvent (e.g. THF, Et₂O) with a strong base (sodium hydride preferred, sodium hexamethyldisilazide, potassium hydride) followed by reaction at
10 reduced temperature (-50 to -20 °C) with n-butyllithium and *N,N,N',N'*-tetramethylethylenediamine followed after a suitable period of time by reaction with anhydrous zinc chloride or magnesium bromide.

- Reaction of the indoline-2-one derivative 6 with either Lawesson's reagent or phosphorous pentasulfide in a suitable organic solvent (pyridine, THF, dioxane,
15 dimethoxyethane, dichloromethane, benzene, toluene, xylene) at a temperature between room temperature and the reflux temperature of the solvent, under an inert atmosphere (nitrogen or argon) provides access to the thiocarbonyl derivative 15. An additive such as sodium hydrogen carbonate may also be useful.

- 34 -


Scheme 6



According to scheme 6 thioamide derivatives 7 may be converted into enamine derivatives 16 (Wrobel, et al, J. Med. Chem., 1989, 2493).

- 5 Thus reaction of thioamide 7 (Pg = H, 2-(trimethylsilyl)-ethoxymethyl, benzyl, etc) with triethyloxonium tetrafluoroborate followed by reaction with a nucleophile (nitromethane, cyanamide, trifluoromethanesulfonamide, Meldrum's acid, etc) followed by removal of the protecting group under appropriate conditions (e.g. tetrabutylammonium fluoride in THF for Pg = 2-(trimethylsilyl)-ethoxymethyl)
- 10 then gives the enamine derivatives 16. Appropriate solvents for the two steps are selected from dichloromethane, THF, dioxane, 1,2-dichloroethane, and the reaction is conducted at a temperature from -78 °C to the boiling point of the solvent under an inert atmosphere (nitrogen or argon).

- 35 -



Scheme 7


According to Scheme 7, treatment of intermediate 7 with an alkylating agent, e.g., methyl iodide, ethyl iodide, 2,4-dinitrofluorobenzene, or 4-nitrofluorobenzene, in the presence of a suitable base (e.g. an amine base such as pyridine, triethylamine or di-iso-propylethylamine or lithium, sodium, potassium or cesium carbonate) in a suitable organic solvent (e.g. DMF, THF, DMSO, dioxane or acetonitrile) at a temperature between -78°C and the boiling point of the solvent, would then afford thioimino ethers 17. Subsequent reaction of intermediates 17 with hydroxylamine or an acid salt of hydroxylamine (e.g. the hydrochloride) in a suitable solvent (for example but not limited to pyridine, methanol, ethanol, iso-propanol, DMF, THF or DMSO and optionally in the presence of an additive such as a tertiary amine base or sodium or potassium acetate) at a temperature between -78°C and the boiling point of the solvent would then afford the *N*-hydroxyamidines 18.

Similarly treatment of intermediates 17 with a carbon nucleophile such as a malonate derivative (e.g., malononitrile, a cyano acetate ester, a nitro acetate ester or a malonate) in the presence of a suitable base (e.g. an amine base such as pyridine, triethylamine or di-iso-propylethylamine or lithium, sodium, potassium or cesium

- 36 -

carbonate) or a Lewis acid (e.g. boron trifluoride etherate, a lead II salt, titanium tetrachloride, a magnesium II salt, or a silver salt) in a solvent compatible with the chosen base or Lewis acid (e.g. DMF, THF, DMSO, dioxane or acetonitrile, chloroform, benzene, toluene or dichloromethane) would then afford the adduct **19**. If
5 the group R3 in adduct **19** is an ester of a carboxylic acid, then it may be decarboxylated directly to give the enamine derivative **20** by treatment with, e.g. sodium iodide in DMSO at a temperature between room temperature and the boiling point of the solvent. Alternatively the ester may be first hydrolysed to the carboxylic acid (by treatment with an aqueous base (e.g. lithium, sodium, or potassium
10 hydroxide) in a suitable solvent (e.g. THF, dioxane acetonitrile, methanol or ethanol), followed by decarboxylation in the presence of an acid (e.g. hydrochloric or sulfuric acid) in a suitable solvent (e.g. acetonitrile, THF, dioxane) to afford the derivatives **20**. Alternatively the xanthate ester of the carboxylic acid may be prepared by
15 reaction with a base such as sodium or potassium hydride in THF, followed by treatment with carbon disulfide. Subsequent reaction with tributyl tin hydride at elevated temperatures in a solvent such as benzene or toluene under an inert nitrogen or argon atmosphere in the presence of a radical initiator such as benzoyl peroxide or azo-bis-*iso*-butyronitrile would then give the product **20**.

- 37 -



Scheme 8

- An alternative strategy for synthesizing the product 18 is illustrated by
- 5 Scheme 8. Thus the bromide 13 (the corresponding chloride, iodide or triflate ester may also be employed) is treated with an alkylating agent, eg methyl iodide, ethyl iodide, 2,4-dinitrofluoro benzene, or 4-nitro fluorobenzene, in the presence of a suitable base (e.g. an amine base such as pyridine, triethylamine or di-iso-propylethylamine or lithium, sodium, potassium or cesium carbonate) in a suitable organic solvent (e.g. DMF, THF, DMSO, dioxane or acetonitrile) at a temperature
- 10 between -78°C and the boiling point of the solvent, would then afford thioimino ethers 21. Subsequent reaction of intermediate 21 with hydroxylamine or an acid salt of hydroxylamine (e.g. the hydrochloride, hydrobromide) in a suitable solvent (for example but not limited to pyridine methanol, ethanol, iso-propanol, DMF, THF or
- 15 DMSO and optionally in the presence of an additive such as a tertiary amine base or sodium or potassium acetate) at a temperature between -78°C and the boiling point of the solvent, would then afford the *N*-hydroxyimide 22. Intermediate 22 could then be protected with a compatible group (e.g. benzyl ether, acyl derivative,


tetrahydropyranyl ether, methoxy methyl ether, silyl ether) to give the derivative 23. Alternately compound 21 could be reacted directly with a protected hydroxylamine derivative (chosen, but not limited to the protecting groups described above) to directly afford derivative 23. Compound 23 may then be reacted with a palladium salt
5 (e.g. tetrakis(triphenylphosphine)palladium(0) or palladium acetate), in a suitable solvent (e.g. THF, dimethoxyethane, acetone, ethanol or toluene) at room temperature under an inert atmosphere (argon, nitrogen). The mixture is then treated with an aryl or heteroaryl boronic acid or boronic acid ester and a base (sodium carbonate, triethylamine, potassium phosphate) in water or fluoride source (cesium fluoride)
10 under anhydrous conditions, and the reaction may then be heated to the boiling point of the solvent. The required product 24 is then isolated and purified by standard means.

Compound 24 may then be de-protected under the conditions prescribed by the nature of the protecting group. For example if the protecting group is a benzyl
15 ether then treatment with boron tribromide or trimethylsilyl iodide in a suitable solvent (dichloromethane for example) would afford the compound 18. Other methods to remove the benzyl ether would involve hydrogenation (hydrogen gas or other hydrogen source such as cyclohexadiene or ammonium formate) in the presence of a palladium catalyst. Solvents suitable for such a process include methanol,
20 ethanol, THF, ethyl acetate and dioxane, at a temperature between room temperature and the boiling point of the solvent. If the protecting group was an acetal derivative (tetrahydropyranyl or methoxymethyl ethers) then hydrolysis could be effected under acidic conditions (hydrochloric acid, sulfuric acid, p-toluene sulfonic acid or acidic ion exchange resin) in a solvent such as methanol, ethanol, THF dioxane or
25 acetonitrile. If the protecting group was an acyl derivative (acetate, or benzoate for example) then hydrolysis could be effected under acidic conditions as described above or under basic conditions (lithium, sodium or potassium hydroxide) in a solvent such as an alcohol, THF dioxane or acetonitrile at a temperature between
30 room temperature and the boiling point of the solvent. If the protecting group was a silyl ether then compound 18 may be prepared by hydrolysing intermediate 24 under

- 39 -

the acidic conditions described above or alternately by exposing compound **24** to a fluoride source (eg potassium fluoride, cesium fluoride or tetra butyl ammonium fluoride) in a solvent such as an alcohol, THF dioxane or acetonitrile at a temperature between room temperature and the boiling point of the solvent. An inert atmosphere of nitrogen or argon may be necessary.

Another method of synthesizing compound **18** would be to convert the protected N-hydroxy amidine **23** into a boronic acid or boronic acid ester (by lithium halogen exchange followed by quench with tri-isopropyl borate, or palladium catalyzed coupling with diboron pinacolate) and then couple this boronic acid or ester derivative with an aryl chloride, bromide, iodide or triflate under a suitable palladium catalysis system as described previously. Subsequent deprotection as described for Scheme 8 would afford the desired compounds **18**.




Scheme 9

According to Scheme 9, treatment of the N-hydroxyamidine **18** under reducing conditions (e.g. catalytic hydrogenation, iron in acetic acid or hydrazine-raney nickel) would then afford intermediate **25**. Solvents suitable for such a process include methanol, ethanol, THF, ethyl acetate and dioxane, at a temperature between room temperature and the boiling point of the solvent. Protection of the secondary

- 40 -

nitrogen (a tertiary butyl carbamate is shown as a non-limiting example) under standard conditions would then give compound 26. Reaction of compound 26 with an electrophilic cyanating agent (e.g. cyanogen bromide, N-cyanobenzotriazole or cyanogen bromide/ 4-dimethylaminopyridine complex) in a suitable solvent (THF
5 acetonitrile or DMF, optionally in the presence of a base such as pyridine or sodium hydride or potassium tert-butoxide) may then afford the desired compound 27. In some cases the cyanation step may occur with concomitant removal of the secondary nitrogen protecting group, if this deprotection does not occur *in-situ* then a further hydrolysis step would be required.

10 An alternate synthesis of compound 27 may follow that of compound 18, Scheme 8, where an N-cyanoamidine bromide 28, prepared from compound 22 adopting a similar strategy to the reactions shown in scheme 9, could be coupled with a suitable functionalised aryl boronic acid or boronic acid ester to give compound 27. In another strategy intermediate 28 may be converted into the corresponding boronic
15 acid or boronic acid ester and coupled in a Suzuki or Suzuki type palladium coupling with a suitable functionalised aryl bromide.



28

20 The compounds of the present invention can be used in the form of salts derived from pharmaceutically or physiologically acceptable acids or bases. These salts include, but are not limited to, the following salts with inorganic acids such as hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid and, as the case may be, such organic acids as acetic acid, oxalic acid, succinic acid, and maleic acid. Other
25 salts include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium in the form of esters, carbamates and other

- 41 -

conventional "pro-drug" forms, which, when administered in such form, convert to the active moiety *in vivo*.

This invention includes pharmaceutical compositions and treatments which comprise administering to a mammal a pharmaceutically effective amount of one or
5 more compounds as described above, or a pharmaceutically acceptable salt thereof, as agonists of the progesterone receptor.

The progesterone receptor agonists of this invention, used alone or in combination, can be utilized in methods of contraception and the treatment and/or prevention of dysfunctional bleeding, uterine leiomyomata, endometriosis; polycystic
10 ovary syndrome, carcinomas and adenocarcinomas of the endometrium, ovary, breast, colon, prostate. Additional uses of the invention include stimulation of food intake.

When the compounds are employed for the above utilities, they may be combined with one or more pharmaceutically acceptable carriers or excipients, for example, solvents, diluents and the like, and may be administered orally in such
15 forms as tablets, capsules, dispersible powders, granules, or suspensions containing, for example, from about 0.05 to 5% of suspending agent, syrups containing, for example, from about 10 to 50% of sugar, and elixirs containing, for example, from about 20 to 50% ethanol, and the like, or parenterally in the form of sterile injectable solutions or suspensions containing from about 0.05 to 5% suspending agent in an
20 isotonic medium. Such pharmaceutical preparations may contain, for example, from about 25 to about 90% of the active ingredient in combination with the carrier, more usually between about 5% and 60% by weight.

The effective dosage of active ingredient employed may vary depending on the particular compound employed, the mode of administration and the severity of the
25 condition being treated. However, in general, satisfactory results are obtained when the compounds of the invention are administered at a daily dosage of from about 0.5 to about 500 mg/kg of animal body weight, preferably given in divided doses two to four times a day, or in a sustained release form. For most large mammals, the total daily dosage is from about 1 to 100 mg, preferably from about 2 to 80 mg. Dosage
30 forms suitable for internal use comprise from about 0.5 to 500 mg of the active

- 42 -

compound in intimate admixture with a solid or liquid pharmaceutically acceptable carrier. This dosage regimen may be adjusted to provide the optimal therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

5 These active compounds may be administered orally as well as by intravenous, intramuscular, or subcutaneous routes. Solid carriers include starch, lactose, dicalcium phosphate, microcrystalline cellulose, sucrose and kaolin, while liquid carriers include sterile water, polyethylene glycols, non-ionic surfactants and edible oils such as corn, peanut and sesame oils, as are appropriate to the nature of the active ingredient and the particular form of administration desired. Adjuvants customarily employed in the preparation of pharmaceutical compositions may be advantageously included, such as flavoring agents, coloring agents, preserving agents, and antioxidants, for example, vitamin E, ascorbic acid, BHT and BHA.

10 The preferred pharmaceutical compositions from the standpoint of ease of preparation and administration are solid compositions, particularly tablets and hard-filled or liquid-filled capsules. Oral administration of the compounds is preferred.

15 These active compounds may also be administered parenterally or intraperitoneally. Solutions or suspensions of these active compounds as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid, polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

20 The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringe ability exists. It must be stable under conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacterial and fungi. The carrier can

25

30

be a solvent or dispersion medium containing, for example, water, ethanol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oil.

This invention is further understood by the following non-limiting examples.

5

EXAMPLE 1

5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indole]-2'(1'H)-thione

Spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one

10 A solution of oxindole (25 g, 0.19 mol) in anhydrous tetrahydrofuran (800 cm³) was cooled to -20°C, then *n*-butyllithium (2.5M in hexanes, 152 cm³, 0.38 mol) was added slowly followed by *N,N,N',N'*-tetramethylethylenediamine (51 cm³, 0.38 mol.). After 15 min. 1,5-diiodopentane (174 g, 0.54 mol) was added slowly and the mixture was allowed to warm to room temperature. After stirring for 16 h. saturated
15 aqueous ammonium chloride solution (1L) and EtOAc (1L) were added. After 15 min., the layers were separated and the aqueous phase was extracted with EtOAc (x2). The combined organic layers were extracted with hydrochloric acid (1N), then washed with brine (500 cm³), dried (MgSO₄), and concentrated to obtain an oil. The oil was triturated with hexane (200 cm³) and benzene (20 cm³). The precipitate was
20 collected and dried *in vacuo* to obtain the subtitled compound (26.3g, 69.6%) as colorless crystals: mp 110-114°C; ¹H NMR (DMSO-d₆) δ 1.67 (m, 10H), 6.84 (d, 1H, *J* = 8 Hz) 6.94 (t, 1H, *J* = 8 Hz), 7.17 (t, 1H, *J* = 8 Hz), 7.44 (d, 1H, *J* = 8 Hz), 10.3 (S, 1H).

25 5'-Bromospiro[cyclohexane-1,3'-[3H] indol]-2'(1'H)-one

To a solution of spiro[cyclohexane-1,3'-[3H]indol]-2' (1'H)-one (17.6 g, 0.09 mol) in acetic acid (300 cm³) was added sodium acetate (8.0 g, 0.1 mol) and bromine (14.6g, 0.091 mol) with stirring. After 30 min. at room temperature, the reaction mixture was partitioned between water and EtOAc. The aqueous phase was extracted
30 twice with EtOAc. The combined organic layers were washed with water, dried

- 44 -

(MgSO₄) and evaporated and the residue was triturated with hexane. The precipitate was collected, and dried *in vacuo* to obtain the subtitled compound (16.5g, 67%) as off-white crystals: mp 196 -199 °C; ¹H NMR (DMSO-d₆) δ 1.62 (m, 10H), 6.8 (d, 1H, J = 6.8 Hz), 7.36 (d, 1H, J = 8.2, 1.8 Hz), 7.58 (dd, 1H, J = 8.2, 1.8 Hz), 10.44 (S, 1H).

5-(3-chlorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-one

A solution of 5'-bromospiro[cyclohexane-1,3'-[3H] indol]-2'(1'H)-one (0.32 g, 1.14 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.14 g, 0.12 mmol) in dimethoxyethane (6 cm³) was stirred under N₂ for 20 min.. To this mixture was then added 3-chlorophenylboronic acid (0.21 g, 1.37 mmol) and sodium carbonate (0.36 g, 3.4 mmol) in water (3 cm³). The solution was brought to reflux for 6 h then cooled to RT, poured into water and extracted with EtOAc (x 3). The combined organic extracts were washed with water, brine, dried (MgSO₄), and evaporated. The residue was purified by column chromatography (SiO₂, ethyl acetate: hexane 1:3) to afford the subtitled compound (0.28 g, 0.89 mmol, 80%) as a yellow solid: mp. 164-165 °C, ¹H NMR (CDCl₃) δ 1.60-1.78 (m, 6H), 1.81-1.99 (m, 4H), 7.04 (d, J = 8.1 Hz, 1H), 7.22-7.47 (m, 4H), 7.53 (s, 1H), 7.61 (s, 1H), 9.28 (br s, 1H); ¹³C-NMR ((CDCl₃) 20.17, 24.12, 31.92 (t), 47.22 (s), 109.21, 121.94, 124.06, 125.50, 125.79, 125.97, 126.38, 128.96 (d), 132.88, 133.59, 135.60, 139.14, 142.17, 182.89 (s); MS (EI) *m/z* 310, 312 (M-H)⁺; Anal. (C₁₉H₁₈ClNO) C, H, N.

To a solution of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (0.63 g, 2.0 mmol) in dry xylene (20 cm³) under nitrogen was added 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (0.89 g, 2.2 mmol) and the mixture heated under reflux. After 72h, the mixture was evaporated and the residue subjected to column chromatography (SiO₂, EtOAc: hexane, gradient elution) to afford a solid which was re-crystallized from di-iso-propylether/ hexane to afford the title compound as yellow crystals (0.17g, 0.51 mmol, 26%): mp. 223 - 227 °C; d (CDCl₃) 1.53 - 1.66 (m, 8H), 1.83 - 2.05 (m, 4H), 2.07 - 2.17 (m, 2H), 7.11 (d, 1H, J = 8.0 Hz) 7.31 - 7.53 (m, 3H), 7.54 (s, 1H), 7.86 (S, 1H), 9.93 (s, 1H, br): MS ((+APCI) *m/z* 328 (M + H)⁺.

EXAMPLE 2

3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol-5'-yl) benzonitrile

To a solution of 5'-bromospiro[cyclohexane-1,3'-[3H] indol]-2'(1'H)-one (1.00 g, 3.57 mmol) in dimethoxyethane (20 cm³) was added

- 5 tetrakis(triphenylphosphine)palladium (0.20 g, 0.17 mmol). After 15 min. 3-formylphenylboronic acid (1.00 g, 6.93 g) was added followed by potassium carbonate (2.90 g, 21 mmol) in water (10 cm³). After 20h at reflux, the mixture was cooled poured into water and extracted with EtOAc (x3). The combined organic extract was washed with saturated brine, dried (MgSO₄) and evaporated. The residue
- 10 was purified by column chromatography (SiO₂, EtOAc: hexane, gradient elution) to afford the title compound (0.66 g, 2.15 mmol, 60%) as a white solid, ¹H NMR (CDCl₃) δ 1.65 – 1.85 (m, 6H), 1.86 - 2.08 (m, 4H), 7.22 (d, 1H, *J* = 8 Hz), 7.48 (dd, 1H, *J* = 8, 2 Hz), 7.61 (t, 1H, *J* = 8 Hz), 7.66 (d, 1H, *J* = 2 Hz), 7.81 - 7.88 (m, 2H), 8.06 (t, 1H, *J* = 2 Hz), 8.30 (s, 1H, br); MS ((+)ESI) *m/z* 306 (M + H)⁺.

15 3-(1',2'-Dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol-5'-yl) benzaldehyde oxime

To a solution of 3-(1',2'-dihydro-2'-oxospirocyclohexane-1,3'-[3H]indol-5'-yl) benzaldehyde (0.59 g, 1.95 mmol) in EtOH: H₂O (10 cm³, 8:2) was added hydroxylamine hydrochloride (0.17 g, 2.5 mmol) and sodium acetate (0.20 g, 2.5 mmol). After 20 min. the mixture was concentrated water was added and the product

20 extracted with EtOAc (x2). The combined organic layers were washed with sat. sodium hydrogen carbonate solution, water, sat. brine, dried (MgSO₄) and evaporated to afford the subtitled oxime (0.63 g, 1.95 mmol, 100%) which was used without further purification, ¹H NMR (CDCl₃) δ 1.60 - 1.84 (m, 6H), 1.85 - 2.00 (m, 4H), 6.86 (d, 1H, *J* = 8 Hz), 7.36 (dd, 1H, *J* = 8, 2 Hz), 7.43 - 7.50 (m, 1H), 7.57 - 7.67 (m, 2H),

25 7.85 (s, 1H, br), 8.25 (s, 1H), 8.68 (s, 1H, br), 8.94 (s, 1H, br); MS ((-)ESI) *m/z* 319 (M - H)⁻.

3-(1',2'-Dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol-5'-yl) benzonitrile

- A solution of 3-(1',2'-Dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol-5'-yl) benzaldehyde oxime (0.48 g, 1.49 mmol) in chloroform (10 cm³) was treated with
- 30 selenium dioxide (0.38 g, 3.50 mmol) and heated under reflux. After 16 h, the

- 46 -

mixture was concentrated and the residue purified by column chromatography (SiO₂, EtOAc: hexane 1:4) and the product re-crystallized from EtOAc-hexane to afford the subtitled compound (0.161 g, 0.53 mmol, 35%) as a white solid: mp. 190 - 191 °C; ¹H NMR (CDCl₃) δ 1.59 - 1.87 (m, 6H), 1.88 - 2.09 (m, 4H), 7.03 (d, 1H, *J* = 8 Hz), 7.42 (dd, 1H, *J* = 8, 2 Hz), 7.54 (t, 1H, *J* = 8 Hz), 7.58 - 7.65 (m, 2H), 7.78 (dt, 1H, *J* = 7, 2 Hz), 7.83 (m, 1H), 8.26 (s, 1H, br); MS ((+) ESI) *m/z* 303 (M + H)⁺.

Reaction of 3-(1',2'-Dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol-5'-yl) benzonitrile and Lawesson's reagent according to the procedure in example 1 gave the title compound: mp. >231 °C (decomp.); ¹H NMR (DMSO-d₆) δ 1.38 - 1.55 (m, 3H), 1.82 - 1.99 (m, 7H), 7.16 (d, 1H, *J* = 8.1 Hz), 7.63 - 7.69 (m, 2H), 7.80 (d, 1H, *J* = 7.7 Hz), 8.01 (d, 1H, *J* = 8 Hz) and 12.76 (s, 1H); MS ((-)APCI) *m/z* 317 [M-H]⁻.

EXAMPLE 3

4-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile

3-(Trimethylstannyl)-2-thiophenecarbonitrile. A solution of 3-bromo-2-thiophenecarbonitrile (0.8 g, 4.3 mmol), tetrakis(triphenylphosphine)palladium(0) (0.25 g, 0.2 mmol) and hexamethylditin (1.4 g, 4.3 mmol) in dimethoxyethane (5 cm³) was heated under reflux for 14 h then cooled to RT. The reaction mixture was absorbed onto florisil and purified by column chromatography (SiO₂, methylene chloride: hexane 1:9) to afford the subtitled compound (1.04 g, 3.8 mmol, 90%) as a clear viscous oil: ¹H NMR (CDCl₃) δ 0.35 (s, 9H), 7.56 (d, *J* = 0.9 Hz, 1H), 7.66 (d, *J* = 0.9 Hz, 1H).

4-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-thiophenecarbonitrile

A solution of the 5'-bromospiro[cyclohexane-1,3'-[3H] indol]-2'(1'H)-one (0.53 g, 1.9 mmol), dichlorobis(triphenylphosphine) palladium(II) (0.1 g, 0.14 mmol) and triphenylarsine (0.14 g, 0.47 mmol) in dimethoxyethane (8 cm³) was stirred under N₂ for 20 minutes. To this mixture was then added 3-(trimethylstannyl)-2-thiophenecarbonitrile (0.64 g, 2.35 mmol). The solution was brought to reflux for 32 h. After cooling to room temperature the reaction mixture was absorbed onto florisil

and purified by column chromatography (SiO₂, ethyl acetate: hexane 2:3) to afford the subtitled compound (0.43 g, 1.39 mmol, 74%) as an off white solid: ¹H NMR (CDCl₃) δ 1.56-2.1 (m, 10H), 6.97 (d, *J* = 8.0 Hz, 1H), 7.39 (dd, *J* = 8.03, 1.45 Hz, 1H), 7.57 (d, *J* = 1.45 Hz, 1H), 7.59 (d, *J* = 1.4 Hz, 1H), 7.84 (d, *J* = 1.4 Hz, 1H), 8.32 (br s, 1H); ¹³C-NMR (CDCl₃) δ 22.07, 26.56, 34.4 (t), 48.13 (s), 110.18 (d), 111.3, 114.75 (s), 122.92, 126.76 (d), 128.44 (s), 137.55 (d), 138.11, 142.71, 144.49, 182.13 (s); MS (EI) *m/z* 307 (M-H)⁺; Anal. (C₁₈H₁₆N₂OS) C, H, N.

A solution of 4-(1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-thiophenecarbonitrile (1.0 g, 3.2 mmol) and Lawesson's Reagent (1.3 g, 3.2 mmol) in o-xylene (20 mL) was heated for two and a half hours. The reaction mixture was washed with distilled water (5x100 mL), dried over MgSO₄, and evaporated. The product was purified by column chromatography (SiO₂, EtOAc:Hexane 1:5) to afford the title compound (0.2 g, 20%) as a pale-yellow solid: m.p. 230-232 °C; ¹H-NMR (DMSO-d₆) δ 12.72 (s, 1H), 8.52 (d, 1H, *J* = 1.5 Hz), 8.36 (d, 1H, *J* = 1.5 Hz), 8.00 (d, 1H, *J* = 1.5 Hz), 7.69 (dd, 1H, *J* = 6.4, 1.8 Hz), 7.10 (d, 1H, *J* = 8.3 Hz), 1.98 - 1.77 (m, 7H), 1.43 - 1.33 (m, 3H); MS (EI) M⁺ @ *m/z* 324.

EXAMPLE 4

3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile

To a solution of 5'-bromospiro [cyclohexane-1,3'-[3H]indol]-2'-(1'-H)-one (11 g, 0.04 mol) in dry tetrahydrofuran (200 cm³) was added sodium hydride (60% dispersion in mineral oil, 1.6 g, 0.04 mol). After 30 min. stirring at room temperature, the mixture was cooled to -78°C and butyl lithium (1.7M in hexanes, 23.2 cm³, 0.04 mol) was added slowly. After 30 min. di-*iso*-propylborate (25 cm³, 0.11 mol) was added and the mixture was allowed to warm to room temperature. After 2 hrs. hydrochloric acid (1N, 500 cm³) and ethylacetate (500 cm³) were added. The aqueous phase was extracted with ethylacetate, then the combined organic layers were washed with water, brine, dried (MgSO₄) and evaporated. The residue was triturated with hexane and the precipitate dried *in vacuo* to obtain (2'-oxo-2, 3-

- 48 -

dihydrospiro[cyclohexane-1, 3'-[3H] indol] -5'-yl) boronic acid (8.3 g, 86%) as an off-white solid that was used without further purification. A sample that was further triturated with ethyl acetate had the following properties: mp. 255-260°C dec.; ¹H NMR (DMSO-d₆) δ 1.50 (m, 2H), 1.73 (m, 8H), 6.82 (d, 1H, *J* = 7.72 Hz) 7.66 (d, 1H, *J* = 7.72 Hz) 7.91 (s, 3H, br), 10.36 (s, 1H); MS ((-)ESI) *m/z* 244 [M-H].

3-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile.

To a solution of 3,5-dibromofluorobenzene in diethyl ether (100 cm³) at -78 °C was added *n*-butyl lithium (2.5 M, 8 cm³, 20 mmol) dropwise. After 30 min. the mixture was treated with DMF (20 cm³) in diethyl ether (10 cm³) and stirring was continued at -78 °C. After 30 min. the mixture was quenched with dilute HCl aq., separated and the aqueous layer was extracted with EtOAc. The combined organic layers were combined, washed with water, brine, dried (MgSO₄) and evaporated to give 3-fluoro-5-bromobenzaldehyde (4.0 g, 19.7 mmol, 100%) as an oil: ¹H NMR (CDCl₃) δ *inter alia* 7.50 - 7.53 (m, 2H), 7.82 (s, 1H) and 9.93 (m, 1H); MS (EI) *m/z* 202, 204 [M⁺].

To a solution of the last cited compound (4.0 g, 19.7 mmol) in ethanol:water (8:2, 50 cm³), was added sodium acetate (1.72 g, 21 mmol) and hydroxylamine hydrochloride (1.45 g, 21 mmol), and the mixture was heated under reflux. After 30 min., the mixture was cooled, evaporated and the residue partitioned between water and EtOAc. The aqueous layer was re-extracted with EtOAc and the combined organic layers were washed with water, saturated sodium hydrogen carbonate solution, brine, dried (MgSO₄) and evaporated to give 3-fluoro-5-bromobenzaldehyde oxime (3.76 g, 17.24 mmol, 87%) which was used without further purification: ¹H NMR (CDCl₃) δ 7.24 - 7.27 (m, 2H), 7.50 (s, 1H), 7.68 (s, 1H) and 8.04 (s, 1H); MS (EI) *m/z* 217 [M⁺].

The above oxime (3.76 g, 17.24 mmol) and copper (II) acetate (370 mg) were dissolved in acetonitrile (100 cm³) under nitrogen and heated under reflux. After 5h, the mixture was evaporated, the residue taken into EtOAc, washed with sulfuric acid (1N), water, brine, dried (MgSO₄) and evaporated to give 3-fluoro-5-

bromobenzonitrile (3.08 g, 15.39 mmol, 89%) which was used without further purification.

The above bromide (3.0 g, 15 mmol) and tetrakis(triphenylphosphine)palladium (0) (0.86 g, 0.75 mmol) were dissolved in dimethoxyethane (130 cm³) under nitrogen. After 15 min. (2'-oxo-2, 3-dihydrospiro[cyclohexane-1, 3'-[3H] indol] -5'-yl) boronic acid (2.82 g, 11.5 mmol) and sodium carbonate (3.1 g, 29.3 mmol) dissolved in water (40 cm³) were added, and the mixture heated under reflux. After 8h the mixture was cooled, poured into water and extracted with EtOAc (x3). The combined organic layers were then washed with water, dried (MgSO₄) and evaporated. The residue was then purified by column chromatography (EtOAc: hexane, gradient elution), and the product recrystallized from methanol to give 3-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile (1.78 g, 5.55 mmol, 48%): mp 199 - 205 °C; ¹H NMR (CDCl₃) δ 1.64 -2.03 (m, 10H), 7.03 (d, 1H, *J* = 8 Hz), 7.31 (dt, 1H, *J* = 7.7 and 1.6 Hz), 7.41 (dd, 1H, *J* = 8, 1.7 Hz), 7.49 (dt, 1H, *J* = 9.6, 2 Hz), 7.58 (d, 1H, *J* = 2 Hz), 7.64 (s, 1H) and 8.37 (s, 1H): MS (EI) *m/z* 320 [M⁺].

To a solution of 3-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile (0.32 g, 1.0 mmol) in xylenes (10 cm³) under nitrogen was added Lawesson's reagent (0.89 g, 2.22 mmol) and the reaction was heated under reflux. After 4h., the mixture was cooled, evaporated and the residue subjected to column chromatography (SiO₂, EtOAc: hexane, gradient elution) to afford (0.143 g, 0.42 mmol, 42%) as a white solid: mp. 236-250 °C; ¹H NMR (CDCl₃) δ 1.54 - 1.66 (m, 3H), 1.86 - 2.18 (m, 7H), 7.16 (d, 1H, *J* = 8.1 Hz), 7.33 - 7.36 (m, 1H), 7.46 - 7.52 (m, 2H), 7.65 (s, 1H), 7.85 (d, 1H, *J* = 1Hz), 10.05 (s, 1H); MS ((+)-APCI) *m/z* 337 [M+H]⁺.

EXAMPLE 5

4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophene thioamide

2'-oxo-2',3'-dihydrospiro [cyclohexane-1,3'-[3H]indol]-5'-yl)boronic acid
5 (2.45 g, 10 mmol), 2-bromo-5-cyano-3-methylthiophene (2.4 g, 12 mmol), potassium
(4 g, 29 mmol), and tetrakis(triphenylphosphine)palladium(0) (0.6 g, 0.5 mmol) in
dimethoxyethane: water: ethanol (130 cm³, 10:2:1) was heated to 80°C for 16 h., then
poured into 1L of water, and extracted with EtOAc. The organic layer was washed
with brine, dried (MgSO₄) and concentrated. The crude product was subjected to
10 column chromatography (SiO₂, EtOAc:hexane, 1:1) to obtain the title compound (0.9
g, 28%): m.p. 200-203°C; ¹H NMR (DMSO-d₆) δ 1.63 (m, 8H), 1.87 (m, 2H), 2.27 (s,
3H), 6.95 (d, 1H, J = 8.13 Hz), 7.34 (dd, 1H, J = 8.13, 1.98 Hz) 7.54 (d, 1H, J = 1.98
Hz), 7.82 (s, 1H) 10.50 (s, 1H); MS ((+)APCI) m/z 323 [M + H]⁺.

A solution of 4-methyl-5-[2'-oxo-2'3'-dihydrospiro[cyclohexane-1,3'-
15 [3H]indol]-5'-yl)-2-thiophenecarbonitrile (0.61 g, 1.9 mmol) and phosphorous
pentasulfide (0.92 g, 2.1 mmol) in dioxane (17 mL) was heated to 85 °C for 30
minutes. The reaction mixture was poured into distilled water, and washed with
aqueous NaHCO₃, distilled water, dried over MgSO₄, and evaporated to dryness. The
residue was purified with column chromatography (2.5% MeOH/CH₂Cl₂) to afford
20 the title compound (0.05g, 8%) as a orange-brown solid: m.p. 244-249 °C; ¹H-NMR
(DMSO-d₆) δ 12.75 (s, 1H), 9.54 (s, 1H), 9.34 (s, 1H), 7.76 (d, 1H, J = 1.5 Hz), 7.58
(s, 1H), 7.45 (dd, 1H, J = 6.4, 1.8 Hz), 7.14 (d, 1H, J = 7.9 Hz), 2.26 (s, 3H), 1.98 -
1.89 (m, 7H), 1.83 - 1.81 (m, 3H); MS ((+)APCI) [M+H]⁺ @ m/z 373.

25 **EXAMPLE 6 - Pharmacology**

The progestational activity for the compounds of the current invention was
evaluated in the in-vitro and in-vivo assays described below. In-vitro potencies lie in
the range 0.01 nM - 10,000 nM, and in-vivo potencies in the range 1 µg/kg to 30
mg/kg.

A. In-vitro biology

The in-vitro biology is determined by (1) competitive Radioligand Binding: using the A-form of the human progesterone receptor with progesterone as the radioligand; (2) co-transfection assay, which provides functional activity expressed as agonist EC50 and Antagonist IC50 values; (3) a T47D cell proliferation, which is a further functional assay which also provides agonist and antagonist data; and (4) T47D cell alkaline phosphatase assay, which is a further functional assay which also provides agonist and antagonist data.

1. hPR Binding assay - This assay is carried out in accordance with: Pathirana, C.; Stein, R.B.; Berger, T.S.; Fenical, W.; Ianiro, T.; Mais, D.E.; Torres, A.; Glodman, M.E., *Nonsteroidal human progesterone receptor modulators from the marine alga cymoplia barbata*, J. Steroid Biochem. Mol. Biol., 1992, **41**, 733-738.

2. PRE-luciferase assay in CV-1 cells

The object of this assay is to determine a compound's progestational or antiprogestational potency based on its effect on PRE-luciferase reporter activity in CV-1 cells co-transfected with human PR and PRE-luciferase plasmids. The materials methods used in the assay are as follows.

a. Growth medium: DMEM (BioWhittaker) containing 10% (v/v) fetal bovine serum (heat inactivated), 0.1 mM MEM non-essential amino acids, 100U/ml penicillin, 100mg/ml streptomycin, and 2 mM GlutaMax (GIBCO, BRL). Experimental medium: DMEM (BioWhittaker), phenol red-free, containing 10% (v/v) charcoal-stripped fetal bovine serum (heat-inactivated), 0.1 mM MEM non-essential amino acids, 100U/ml penicillin, 100mg/ml streptomycin, and 2 mM GlutaMax (GIBCO, BRL).

b. Cell culture, transfection, treatment, and luciferase assay

Stock CV-1 cells are maintained in growth medium. Co-transfection is done using 1.2×10^7 cells, 5 mg pLEM plasmid with hPR-B inserted at SphI and BamHI sites, 10 mg pGL3 plasmid with two PREs upstream of the

- 52 -

luciferase sequence, and 50 mg sonicated calf thymus DNA as carrier DNA in 250 ml. Electroporation is carried out at 260 V and 1,000 mF in a Biorad Gene Pulser II. After electroporation, cells are resuspended in growth medium and plated in 96-well plate at 40,000 cells/well in 200 μ l. Following overnight incubation, the medium is changed to experimental medium. Cells are then treated with reference or test compounds in experimental medium. Compounds are tested for antiprogestational activity in the presence of 3 nM progesterone. Twenty-four hr. after treatment, the medium is discarded, cells are washed three times with D-PBS (GIBCO, BRL). Fifty μ l of cell lysis buffer (Promega, Madison, WI) is added to each well and the plates are shaken for 15 min in a Titer Plate Shaker (Lab Line Instrument, Inc.). Luciferase activity is measured using luciferase reagents from Promega.

c. Analysis of Results:

Each treatment consists of at least 4 replicates. Log transformed data are used for analysis of variance and nonlinear dose response curve fitting for both agonist and antagonist modes. Huber weighting is used to downweight the effects of outliers. EC_{50} or IC_{50} values are calculated from the retransformed values. JMP software (SAS Institute, Inc.) is used for both one-way analysis of variance and non-linear response analyses.

d. Reference Compounds:

Progesterone and trimegestone are reference progestins and RU486 is the reference antiprogesterin. All reference compounds are run in full dose-response curves and the EC_{50} or IC_{50} values are calculated.

Table 1. Estimated EC₅₀, standard error (SE), and 95% confidence intervals (CI) for reference progestins from three individual studies

5	Compound	Exp.	EC ₅₀	SE	95% CI	
			(nM)		lower	upper
	Progesterone	1	0.616	0.026	0.509	0.746
		2	0.402	0.019	0.323	0.501
		3	0.486	0.028	0.371	0.637
10	Trimegestone	1	0.0075	0.0002	0.0066	0.0085
		2	0.0081	0.0003	0.0070	0.0094
		3	0.0067	0.0003	0.0055	0.0082

Table 2. Estimated IC₅₀, standard error (SE), and 95% confident interval (CI) for the antiprogesterin, RU486 from three individual studies

15	Compound	Exp.	IC ₅₀	SE	95% CI	
			(nM)		lower	upper
20	RU486	1	0.028	0.002	0.019	0.042
		2	0.037	0.002	0.029	0.048
		3	0.019	0.001	0.013	0.027

Progestational activity: Compounds that increase PRE-luciferase activity significantly ($p < 0.05$) compared to vehicle control are considered active.

25 Antiprogestational activity: Compounds that decrease 3 nM progesterone induced PRE-luciferase activity significantly ($p < 0.05$)

EC₅₀: Concentration of a compound that gives half-maximal increase PRE-luciferase activity (default-nM) with SE.

30 IC₅₀: Concentration of a compound that gives half-maximal decrease in 3 nM progesterone induced PRE-luciferase activity (default-nM) with SE.

- 54 -

3. T47D cell proliferation assay

The objective of this assay is the determination of progestational and antiprogesterational potency by using a cell proliferation assay in T47D cells. A compound's effect on DNA synthesis in T47D cells is measured. The materials and methods used in this assay are as follows.

a. Growth medium: DMEM:F12 (1:1)

(GIBCO, BRL) supplemented with 10% (v/v) fetal bovine serum (not heat-inactivated), 100U/ml penicillin, 100mg/ml streptomycin, and 2 mM GlutaMax (GIBCO, BRL).

b. Treatment medium: Minimum Essential

Medium (MEM) (#51200-038 GIBCO, BRL) phenol red-free supplemented with 0.5% charcoal stripped fetal bovine serum, 100U/ml penicillin, 200 mg/ml streptomycin, and 2 mM GlutaMax (GIBCO, BRL).

c. Cell culture

Stock T47 D cells are maintained in growth medium. For BrdU incorporation assay, cells are plated in 96-well plates (Falcon, Becton Dickinson Labware) at 10,000 cells/well in growth medium. After overnight incubation, the medium is changed to treatment medium and cells are cultured for an additional 24 hr before treatment. Stock compounds are dissolved in appropriate vehicle (100% ethanol or 50% ethanol/50% DMSO), subsequently diluted in treatment medium and added to the cells. Progesterin and antiprogesterin reference compounds are run in full dose-response curves. The final concentration of vehicle is 0.1%. In control wells, cells receive vehicle only. Antiprogesterins are tested in the presence of 0.03 nM trimegestone, the reference progesterin agonist. Twenty-four hours after treatment, the medium is discarded and cells are labeled with 10 mM BrdU (Amersham Life Science, Arlington Heights, IL) in treatment medium for 4 hr.

d. Cell Proliferation Assay

At the end of BrdU labeling, the medium is removed and BrdU incorporation is measured using a cell proliferation ELISA kit (#RPN 250, Amersham Life Science) according to manufacturer's instructions.

- 55 -

Briefly, cells are fixed in an ethanol containing fixative for 30 min, followed by incubation in a blocking buffer for 30 min to reduce background. Peroxidase-labeled anti-BrdU antibody is added to the wells and incubated for 60 min. The cells are rinsed three times with PBS and incubated with 3,3',5,5'-tetramethylbenzidine (TMB) substrate for 10-20 min depending upon the potency of tested compounds. Then 25 μ l of 1 M sulfuric acid is added to each well to stop color reaction and optical density is read in a plate reader at 450 nm within 5 min.

e. Analysis of Results:

Square root-transformed data are used for analysis of variance and nonlinear dose response curve fitting for both agonist and antagonist modes. Huber weighting is used to downweight the effects of outliers. EC₅₀ or IC₅₀ values are calculated from the retransformed values. JMP software (SAS Institute, Inc.) is used for both one-way analysis of variance and non-linear dose response analyses in both single dose and dose response studies.

f. Reference Compounds:

Trimegestone and medroxyprogesterone acetate (MPA) are reference progestins and RU486 is the reference antiprogesterin. All reference compounds are run in full dose-response curves and the EC₅₀ or IC₅₀ values are calculated.

Table 3. Estimated EC₅₀, standard error (SE), and 95% confidence intervals (CI) for individual studies

Compound	Exp	EC ₅₀ (nM)	SE	95% CI	
				lower	upper
Trimegestone	1	0.017	0.003	0.007	0.040
	2	0.014	0.001	0.011	0.017
	3	0.019	0.001	0.016	0.024
MPA	1	0.019	0.001	0.013	0.027
	2	0.017	0.001	0.011	0.024

Table 4. Estimated IC₅₀, standard error, and 95% confident interval for the antiprogesterin, RU486

5	Compound	Exp	IC ₅₀ (nM)	SE	95% CI	
					lower	upper
	RU486	1	0.011	0.001	0.008	0.014
		2	0.016	0.001	0.014	0.020
		3	0.018	0.001	0.014	0.022

- 10 EC₅₀: Concentration of a compound that gives half-maximal increase in BrdU incorporation with SE; IC₅₀: Concentration of a compound that gives half-maximal decrease in 0.1 trimegestone induced BrdU incorporation with SE

4. T47D cell alkaline phosphatase assay

The purpose of this assay is to identify progestins or

- 15 antiprogestins by determining a compound's effect on alkaline phosphatase activity in T47D cells. The materials and methods used in this assay are as follows.

a. Culture medium: DMEM:F12 (1:1) (GIBCO, BRL) supplemented with 5% (v/v) charcoal stripped fetal bovine serum (not heat-inactivated), 100U/ml penicillin, 100 µg/ml streptomycin, and 2 mM GlutaMax (GIBCO, BRL).

- 20 b. Alkaline phosphatase assay buffer:

I. 0.1 M Tris-HCl, pH 9.8, containing 0.2% Triton X-100; II. 0.1 M Tris-HCl, pH 9.8 containing 4 mM p-nitrophenyl phosphate (Sigma).

c. Cell Culture and Treatment:

Frozen T47D cells were thawed in a 37°C water

- 25 bath and diluted to 280,000 cells/ml in culture medium. To each well in a 96-well plate (Falcon, Becton Dickinson Labware), 180 µl of diluted cell suspension was added.

Twenty µl of reference or test compounds diluted in the culture medium was then added to each well. When testing for progestin antagonist activity, reference antiprogestins or test compounds were added in the presence of 1 nM progesterone.

- 30 The cells were incubated at 37°C in a 5% CO₂/humidified atmosphere for 24 hr.

- 57 -

d. Alkaline Phosphatase Enzyme Assay:

At the end of treatment, the medium was removed from the plate and fifty μ l of assay buffer I was added to each well. The plates were shaken in a titer plate shaker for 15 min. Then 150 μ l of assay buffer II was added to each well. Optical density measurements were taken at 5 min intervals for 30 min at a test wavelength of 405 nM.

e. Analysis of Results: Analysis of dose-response data

For reference and test compounds, a dose response curve is generated for dose (X-axis) vs. the rate of enzyme reaction (slope) (Y-axis). Square root-transformed data are used for analysis of variance and nonlinear dose response curve fitting for both agonist and antagonist modes. Huber weighting is used to downweight the effects of outliers. EC_{50} or IC_{50} values are calculated from the retransformed values. JMP software (SAS Institute, Inc.) is used for both one-way analysis of variance and non-linear dose response analyses in both single dose and dose response studies.

f. Reference Compounds:

Progesterone and trimegestone are reference progestins and RU486 is the reference antiprogestin. All reference compounds are run in full dose response curves and the EC_{50} or IC_{50} values are calculated.

Table 5. Estimated EC_{50} , standard error (SE), and 95% confidence intervals (CI) for reference progestins from three independent experiments

Compound	Exp.	EC_{50} (nM)	SE	95% CI	
				lower	upper
Progesterone	1	0.839	0.030	0.706	0.996
	2	0.639	0.006	0.611	0.669
	3	1.286	0.029	1.158	1.429
Trimegestone	1	0.084	0.002	0.076	0.091
	2	0.076	0.001	0.072	0.080
	3	0.160	0.004	0.141	0.181

Table 6. Estimated IC₅₀, standard error, and 95% confident interval for the reference antiprogesterin RU486 from three independent experiments

5	Compound	Exp	IC 50 (nM)	SE	95% CI	
					lower	upper
	RU486	1	0.103	0.002	0.092	0.115
		2	0.120	0.001	0.115	0.126
		3	0.094	0.007	0.066	0.134

10

B. In-vivo Biology

The primary in-vivo assay is the rat decidualization model which may be used to determine progestational effects of both agonists and antagonists. The secondary
 15 in-vivo assay is the rat ovulation inhibition model which is under development and hence the protocol is un-available.

1. Rat decidualization assay: The objective of this procedure is used to evaluate the effect of progestins and antiprogestins on rat uterine decidualization and compare the relative potencies of various test compounds. The
 20 materials and methods used in this assay are as follows.

a. Methods: Test compounds are dissolved in 100% ethanol and mixed with corn oil (vehicle). Stock solutions of the test compounds in oil (MazolaTM) are then prepared by heating (~80 °C) the mixture to evaporate ethanol. Test compounds are subsequently diluted with 100% corn oil or 10%
 25 ethanol in corn oil prior to the treatment of animals. No difference in decidual response was found when these two vehicles were compared.

b. Animals (RACUC protocol #5002)

Ovariectomized mature female Sprague-Dawley rats (~60-day old and 230g) are obtained from Taconic (Taconic Farms, NY) following surgery. Ovariectomy is
 30 performed at least 10 days prior to treatment to reduce circulating sex steroids.

- 59 -

Animals are housed under 12 hr light/dark cycle and given standard rat chow and water *ad libitum*.

c. Treatment

Rats are weighed and randomly assigned to groups of 4
5 or 5 before treatment. Test compounds in 0.2 ml vehicle are administered by
subcutaneous injection in the nape of the neck or by gavage using 0.5 ml. The
animals are treated once daily for seven days. For testing antiprogestins, animals are
given the test compounds and a EC₅₀ dose of progesterone (5.6 mg/kg) during the
first three days of treatment. Following decidual stimulation, animals continue to
10 receive progesterone until necropsy four days later.

d. Dosing

Doses are prepared based upon mg/kg mean group body
weight. In all studies, a control group receiving vehicle is included. Determination of
dose-response curves is carried out using doses with half log increases (e.g. 0.1, 0.3,
15 1.0, 3.0 mg/kg...).

e. Decidual induction

Approximately 24 hr after the third injection, decidualization
is induced in one of the uterine horns by scratching the antimesometrial luminal
epithelium with a blunt 21 G needle. The contralateral horn is not scratched and
20 serves as an unstimulated control. Approximately 24 hr following the final treatment,
rats are sacrificed by CO₂ asphyxiation and body weight measured. Uteri are
removed and trimmed of fat. Decidualized (D-horn) and control (C-horn) uterine
horns are weighed separately.

f. Analysis of Results:

25 The increase in weight of the decidualized uterine horn is
calculated by D-horn/C-horn and logarithmic transformation is used to maximize
normality and homogeneity of variance. The Huber M-estimator is used to down
weight the outlying transformed observations for both dose-response curve fitting and
one-way analysis of variance. JMP software (SAS Institute, Inc.) is used for both one-
30 way ANOVA and non-linear dose-response analyses.

- 60 -

g. Reference Compounds:

All progestin reference compounds were run in full dose-response curves and the EC₅₀ for uterine wet weight were calculated.

5 **Table 7. Estimated EC₅₀, standard error (SE), and 95% confidence intervals for individual studies**

	Compound	Exp	EC ₅₀	SE	95% CI	
			(mg/kg, s.c.)		lower	upper
10	Progesterone	1	5.50	0.77	4.21	7.20
		2	6.21	1.12	4.41	8.76
15	3-Ketodesogestrel	1	0.11	0.02	0.07	0.16
		2	0.10	0.05	0.11	0.25
		3	0.06	0.03	0.03	0.14
20	Levonorgestrel	1	0.08	0.03	0.04	0.16
		2	0.12	0.02	0.09	0.17
		3	0.09	0.02	0.06	0.13
		4	0.09	0.02	0.06	0.14
25	MPA	1	0.42	0.03	0.29	0.60
		2	0.39	0.05	0.22	0.67
		3	0.39	0.04	0.25	0.61

Table 8. Estimated average EC₅₀, standard error, and 95% confidence intervals for dose-response curves of 3 reference compounds

5	Compound	EC ₅₀	SE	95% CI	
		(mg/kg, s.c.)		lower	upper
	Progesterone	5.62	0.62	4.55	7.00
	3-Ketodesogestrel	0.10	0.02	0.07	0.14
	Levonorgestrel	0.10	0.01	0.08	0.12

10 Table 9. Estimated IC₅₀, standard error, and 95% confident interval for the antiprogesterin, RU 486

15	Compound	Exp.	IC ₅₀	SE	95% CI	
			(mg/kg, p.o.)		lower	upper
	RU 486	1	0.21	0.07	0.05	0.96
		2	0.14	0.02	0.08	0.27

Concentration: Compound concentration in assay(default-mg/kg body weight)

Route of administration: Route the compound is administered to the animals

20 Body weight: Mean total animal body weight (default-kg)

D-horn: Wet weight of decidualized uterine horn (default-mg)

C-horn: Wet weight of control uterine horn (default-mg)

Decidual response: [(D-C)/C]x100%

25 Progesterational activity: Compounds that induce decidualization significantly (p<0.05) compared to vehicle control are considered active

Antiprogesterational activity: Compounds that decrease EC₅₀ progesterone induced decidualization significantly (p<0.05)

EC₅₀ for uterine weight: Concentration of compound that gives half-maximal increase in decidual response (default-mg/kg)

30 IC₅₀ for uterine weight: Concentration of compound that gives half-maximal decrease in EC₅₀ progesterone induced decidual response (default-mg/kg)

Table 10. Data for Representative Compounds

Example #	Ki/nM	CV-1 EC50/nM	Ovulation inhibition IC100 mg/kg
5		0.3	
3	0.1	0.2	
1	0.2	0.8	
4	0.06	0.1	0.1

5

Example 7**5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile**

- 5-(2'-Oxo-2',3'-dihydrospiro[cyclopentane-1,3'-[3H]indol]-5'-yl)-2-cyanopyrrole: A solution of 5'-bromospiro[cyclopentane-1,3'-[3H] indol]-2'(1'H)-one (2.0 g, 7.5 mmol) and tetrakis(triphenylphosphine)palladium(0) (430 mg, 0.3 mmol) in ethylene glycol dimethyl ether (50 mL) was stirred under a flow of nitrogen for 15 min. To the solution was added sequentially 1-t-butoxycarbonylpyrrole-2-boronic acid (2.1 g, 9.7 mmol) and potassium carbonate (2.4 g, 17 mmol) in water (10 mL). The mixture was heated to 80 °C for 3 h and allowed to cool. The reaction mixture was poured into water (50 mL) and extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, washed with brine (30 mL) and dried over magnesium sulfate. The solution was filtered and concentrated *in vacuo*. Crystallization from 20% ethyl acetate/hexane gave 2-(1',2'-dihydro-2'-oxospiro[cyclopentane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-1-carboxylic acid, tert-butyl ester (2.2 g, 83%) as a white powder, mp 179-180.5 °C. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 1.30 (s, 9H), 1.75-1.98 (m, 8 H), 6.16 (dd, 1 H, *J* = 1.8, 3.3 Hz), 6.22 ('t', 1 H, *J* = 3.3, 3.3 Hz), 6.79 (d, 1 H, *J* = 7.9 Hz), 7.08 (dd, 1 H, *J* = 1.8, 7.9 Hz), 7.14

- 63 -

('d', 1 H, $J = 1.5$ Hz), 7.28 (dd, $J = 1.9, 3.3$ Hz), 10.30 (s, 1 H). MS (EI) m/z 352 [M^+]. Anal. Calcd for $C_{21}H_{24}N_2O_3$: C, 71.57; H, 6.86; N, 7.95. Found: C, 71.08; H, 6.83; N, 7.74.

To a solution of 2-(1',2'-dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-1-carboxylic acid, tert-butyl ester (2.2 g, 6.0 mmol) in THF (anhydrous, 25 mL) was added at -78°C chlorosulfonyl isocyanate (0.63 mL, 7.0 mmol). After 90 min, dimethylformamide (11 mL, 140 mmol) was added and the reaction was allowed to warm to room temperature. The reaction mixture was poured into water (50 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layers were combined, washed with brine (50 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification via flash column chromatography on silica gel (30% ethyl acetate/hexane) gave 5-(2'-oxo-2',3'-dihydrospiro[cyclopentane-1,3'-[3H]indol]-5'-yl)-2-cyanopyrrole-1-carboxylic acid, tert-butyl ester (1.7 g, 75%) as white crystals, mp $167-9^\circ\text{C}$. ^1H NMR (DMSO- d_6 , 400 MHz) δ 1.34 (s, 9H), 1.75-1.98 (m, 8 H), 6.39 (d, 1 H, $J = 3.7$ Hz), 6.84 (d, 1 H, $J = 7.9$ Hz), 7.17 (dd, 1 H, $J = 1.8, 7.9$ Hz), 7.28 ('t', 2 H), 10.41 (s, 1 H). MS (ESI) m/z 376 [$M-H$] $^-$. Anal. Calcd. for $C_{22}H_{23}N_3O_3$: C, 70.01; H, 6.14; N, 11.13. Found: C, 69.67; H, 6.38; N, 11.04.

5-(2'-Oxo-2',3'-dihydrospiro[cyclopentane-1,3'-[3H]indol]-5'-yl)-2-cyanopyrrole-1-carboxylic acid, tert-butyl ester (1 g, 2.7 mmol) was placed in a 25 mL round bottomed flask stoppered with a rubber septum and equipped with nitrogen inlet and a needle to allow gaseous outflow. A vigorous flow of nitrogen was maintained as the flask was placed in an oil bath and heated to 165°C . After 20 min at this temperature, the flask was removed from the oil bath and allowed to cool. Crystallization from ethyl ether gave the title compound (600 mg, 79%) as a yellow powder, mp $285-286^\circ\text{C}$. ^1H NMR (DMSO- d_6 , 400 MHz) δ 1.75-2.03 (m, 8 H), 6.60 (dd, 1 H, $J = 2.4, 3.7$ Hz), 6.84 (d, 1 H, $J = 8.1$ Hz), 6.94 (dd, 1 H, $J = 2.4, 3.7$ Hz), 7.52 (dd, 1 H, $J = 1.8, 8.1$ Hz), 7.60 (d, 1 H, $J = 1.8$ Hz), 10.38 (s, 1 H), 12.45 (s, 1 H). MS (ESI) m/z 276 [$M-H$] $^-$. Anal. Calcd. For $C_{17}H_{15}N_3O$: C, 73.63; H, 5.45; N, 15.15. Found: C, 73.24; H, 5.34; N, 14.96.

- 64 -

To 5-(1,2-Dihydro-2-oxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile (0.18 g, 0.7 mmol, 1 eq) in *p*-xylene (20 mL) was added Lawesson's reagent (0.14 g, 0.36 mmol, 0.5 eq) and the reaction was heated to reflux for 1 hour. The reaction was cooled to room temperature and adsorbed onto silica gel. Purification by flash column chromatography (20% ethyl acetate/hexane) on silica gel gave the product as an orange powder. Further purification by HPLC gave the title compound as a green solid (0.144 g, 70%), mp 275-276 °C (dec.). ¹H NMR (*d*₆-DMSO, 300 MHz) δ 1.81-2.16 (m, 8 H), 6.69 (dd, 1 H, *J* = 2.3, 3.7 Hz), 6.98 (dd, 1 H, *J* = 1.8, 3.7 Hz), 7.04 (d, 1 H, *J* = 8.2 Hz), 7.63 (dd, 1 H, *J* = 1.6, 8.2 Hz), 7.72 (d, 1 H, *J* = 1.3 Hz), 12.57 (s, 1 H), 12.65 (s, 1 H). MS (ESI) [*M*-H]⁻ = 292. Anal. Calculated (calcd.) for C₁₇H₁₅N₃S: C, 69.6; H, 5.15; N, 14.32. Found: C, 69; H, 5.31; N, 13.81.

Example 8

15 **5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile**

To a solution of 5'-bromo-spiro[cyclohexane-1,3'-indolin]-2'-one (3.4 g, 12 mmol) in 1,2-DME (100 mL) under a nitrogen atmosphere was added tetrakis(triphenylphosphine)palladium(0) (70 mg, 5 mol%). After 15 min, 2-borono-1H-pyrrole-1-carboxylic acid, 1-tert butyl ester (1.3 eq, 3.31 g, 15.6 mmol) and a solution of K₂CO₃ (2.3 eq, 3.83 g, 27.6 mmol) in water (5 mL) were added sequentially. The solution was heated to 80 °C for 3 h and allowed to cool. The reaction mixture was poured into water (200 mL) and extracted with EtOAc (2 x 100 mL). The organic layers were combined, washed with brine (150 mL) and dried over MgSO₄. The solution was filtered, concentrated *in vacuo*, and the residue was purified by flash column chromatography on silica gel (eluting with 30% EtOAc/hexane) to give 2-(1',2'-dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-1-carboxylic acid, tert-butyl ester (3.4 g, 76%) as a white powder, mp 177 °C. ¹H NMR (CDCl₃, 300 MHz) δ 1.38 (s, 9 H), 1.59-1.93 (m, 10 H), 6.18 (m, 1 H), 6.23 (t, 1H, 3 Hz), 6.91 (d, 1 H, *J*=8 Hz), 7.21 (d, 1 H, *J*=8 Hz), 7.34 (m, 1 H),

- 65 -

7.44 (s, 1 H), 8.33 (br s, 1 H, D₂Oex). MS ((+)-APCI) m/z 367 [(M+H)⁺]. Anal. Calcd for C₂₂H₂₆N₂O₃: C, 72.11; H, 7.15; N, 7.64. Found: C, 71.7; H, 7.16; N, 7.5.

To a solution of 2-(1',2'-dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-1-carboxylic acid, tert-butyl ester (0.75 g, 2 mmol) in THF
5 (anhydrous, 20 mL) at -78 °C was added chlorosulfonyl isocyanate (1.15 eq, 0.23 mL, 2.3 mmol). After 90 min, DMF (20 eq, 3.6 mL, 46 mmol) was added and the reaction was allowed to warm to room temperature. The reaction mixture was poured into water (50 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layers were combined, washed with brine (50 mL), dried over magnesium sulfate, filtered and
10 concentrated *in vacuo*. Purification via flash column chromatography on silica gel (30% ethyl acetate/hexane) gave 5-(2'-oxo-2',3'-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-cyanopyrrole-1-carboxylic acid, tert-butyl ester (0.5 g, 63%) as an oil which crystallized from acetone to give white crystals, mp 156 °C. ¹H NMR (*d*₆-DMSO, 400 MHz) δ 1.32 (s, 9H), 1.50 (m, 3 H), 1.60-1.70 (m, 5 H), 1.75-1.85 (m, 2
15 H), 6.38 (d, 1 H, *J* = 3.7 Hz), 6.87 (d, 1 H, *J* = 7.9 Hz), 7.18 (dd, 1 H, *J* = 1.5, 7.9 Hz), 7.27 (d, 1 H, *J* = 3.7 Hz), 7.48 (d, 1 H, *J* = 1.8 Hz), 10.42 (bs, 1 H). MS (EI) m/z 391 (M⁺). Anal. Calcd for C₂₃H₂₅N₃O₃: C, 70.57; H, 6.44; N, 10.73. Found: C, 69.82; H, 6.46; N, 10.43.

To a solution of 2-Cyano-5-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-
20 [3H]indol]-5-yl)-1H-pyrrole-1-carboxylic acid, tert-butyl ester (0.7 g, 1.8 mmol, 1 eq) in toluene (70 mL) was added Lawesson's reagent (0.47 g, 1.1 mmol, 0.65 eq) and the reaction was heated to reflux for 1 hour. The reaction was cooled to room temperature, poured into water (100 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layers were combined, washed with brine (50 mL), dried over
25 magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (20-30% ethyl acetate/hexane) on silica gel gave title compound as a yellow solid (0.7 g, 96 %). ¹H NMR (*d*₆-DMSO, 500 MHz) δ 1.30-1.98 (m, 19 H), 6.45 (d, 1 H, *J* = 3.7 Hz), 7.09 (d, 1 H, *J* = 7.9 Hz), 7.31-7.34 (m, 2 H), 7.81 (d, 1 H, *J* = 1.4 Hz), 12.74 (s, 1 H). MS (ESI) [M-H]⁻ = 406. Anal. calcd. for C₂₃H₂₅N₃O₂S: C, 67.79; H, 6.18; N, 10.31. Found: C, 67.86; H, 5.99; N, 10.25.
30

Example 9**5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-H-pyrrole-2-carbonitrile**

5 To a solution of 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile (0.5 g, 1.2 mmol, 1 eq) in THF (5 mL) was added NaOEt (0.25 g, 3.6 mmol, 3 eq) in EtOH (5 mL) and the reaction was heated to 80°C for 24 h. The solvents were removed *in vacuo* and the residue partitioned between ethyl acetate (50 mL) and water (50 mL). The layers were
10 separated and the aqueous layer was extracted with ethyl acetate (50 mL). The organic layers were combined, washed with brine (50 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (30% ethyl acetate/hexane) on silica gel gave the title compound (0.27g, 68%) as a yellow powder. ¹H NMR (*d*₆-DMSO, 500 MHz) δ 1.32-1.99 (m, 10 H), 6.71 (d, 1 H, *J* = 3.7 Hz), 7.00 (d, 1 H, *J* = 3.7 Hz), 7.09 (d, 1 H, *J* = 8.4 Hz), 7.70 (dd, 1 H, *J* = 1.6, 8.4 Hz), 8.05 (d, 1 H, *J* = 1.1 Hz), 12.67 (s, 1 H), 12.73 (s, 1 H). MS (ESI) [M-H]⁻ = 306. Anal. calcd. for C₁₈H₁₇N₃S: C, 70.33; H, 5.57; N, 13.67. Found: C, 69.64; H, 5.79; N, 13.04.

20

Example 10**5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile**

To a solution of 5-(2'-oxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile (0.55 g, 1.8 mmol, 1 eq) in toluene (50 mL) was added
25 Lawesson's reagent (0.47 g, 1.1 mmol, 0.65 eq) and the reaction was heated to 80 °C for 1 hour. The reaction was cooled to room temperature, poured into water (100 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layers were combined, washed with brine (50 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography on silica gel gave the product
30 as a white solid (0.32 g, 55 %). ¹H NMR (*d*₆-DMSO, 500 MHz) δ 1.36-1.99 (m, 10

- 67 -

H), 3.7 (s, 3 H), 6.35 (d, 1 H, $J = 4.2$ Hz), 7.05 (d, 1 H, $J = 4.2$ Hz), 7.16 (d, 1 H, $J = 7.9$ Hz), 7.44 (dd, 1 H, $J = 1.6, 8.1$ Hz), 7.83 (d, 1 H, $J = 1.6$ Hz), 12.75 (s, 1 H). MS (ESI) $[M-H]^+ = 320$. Anal. calcd. for $C_{19}H_{19}N_3S$: C, 70.99; H, 5.96; N, 13.07. Found: C, 68.69; H, 5.36; N, 12.27.

5

Example 11

5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3-thiophenecarbonitrile

5-Bromo-2-thiophenecarbonitrile: A mixture of 5-bromo-2-

10 thiophenecarboxaldehyde (96.0g, 500 mmol), hydroxylamine hydrochloride (111.9 g, 500 mmol), pyridine (500 mL), and ethanol (500 mL) was heated under nitrogen at reflux for two hours. The reaction mixture was cooled to ambient temperature and concentrated *in vacuo* to give an oil. The crude product was triturated twice with ice water and the solid obtained was collected on a filter. A mixture of a portion of the
15 above solid (44.31 g, 215 mmol), copper (II) acetate monohydrate (4.2 g, 21 mmol) in acetonitrile (1.4L) was heated at reflux for three hours. The solvent was removed *in vacuo* and the residue was dissolved in ethyl acetate. The solution was washed with 5% aqueous sulfuric acid (2X30 mL), water (2X30 mL), brine (20 mL), and dried ($MgSO_4$). The solvent was removed *in vacuo* and the residue was dissolved in a
20 minimum amount of chloroform (1L) and allowed to crystallize. The crystal obtained was collected on a filter and the filtrate was concentrated and purified by a chromatography (silica gel, chloroform) to give the subtitled compound as an off-white solid (31.5g combined, 58%). IR (film) cm^{-1} 2200. 1H -NMR ($CDCl_3$) δ 7.39-7.38 (d, 1H, $J = 4.1$ Hz), 7.10 (d, 1H, $J = 4.0$ Hz); MS (EI) m/z 187 (M^+ , 98%)
25 189(M^+ , 100%).

5-(1,2-dihydro-2-oxospiro[cyclopentane-1,3-[3H]indole]-5-yl)-3-thiophenecarbonitrile was prepared according to the procedure for Example 5 using 5-bromo-2-thiophenecarbonitrile and (2'-oxo-2',3'-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl) boronic acid: mp. 225-228°C; 1H NMR ($DMSO-d_6$) δ 1.63 (m, 8H),
30 1.90 (m, 2H) 6.91 (d, 1H, $J = 8.13$ Hz), 7.55 (dd, 1H, $J = 8.13, 1.76$ Hz), 7.60 (d, 1H,

- 68 -

$J = 4.17$ Hz), 7.75 (d, 1H, $J = 1.76$ Hz), 7.93 (d, 1H, $J = 4.17$ Hz), 10.51 (s, 1H); MS ((+)APCI) m/z 309 [M + H]⁺.

A solution of 5-(1,2-dihydro-2-oxospiro[cyclopentane-1,3-[3H]indole]-5-yl)-3-thiophenecarbonitrile (0.66 g, 2.4 mmol), and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (0.97 g, 2.4 mmol) in toluene (250 ml) was stirred at 80°C for 2 hours. The solution was concentrated in vacuo. The residue was extracted with ethylacetate, the ethylacetate solution was washed with water, dried over magnesium sulfate, and concentrated. The residue was purified by column chromatography (silica gel, ethylacetate, hexane 20/80) to afford the title compound, m.p. 269-272 °C (0.24 g, 32%). ¹H-NMR (DMSO-d₆) δ 2.09 (m, 8H), 7.05 (d, $J = 8.1$ Hz, 1H), 7.55 (dd, $J = 8.1, 1.7$ Hz, 1H), 7.7 (d, $J = 1.7$ Hz, 1H), 7.95 (d, $J = 1.3$ Hz, 1H), 8.49 (d, $J = 1.3$ Hz, 1H), 8.49 (d, $J = 1.3$ Hz, 1H), 12.68 (s, 1H); MS (EI NEG) m/z 309 (M-H)⁻.

15

Example 12**5-(1,2-Dihydro - thioxospiro (cyclopentane - 1,3-[3 H]indol) - 5-yl) -2-thiophenecarbonitrile**

The title compound was prepared from 5-(1,2-dihydro-oxospiro(cyclopentane - 1,3- [3 H] indol)-5 yl)-2-thiophenecarbonitrile (2 g, 6.8 mmol) and Lawesson's reagent (3.32 g, 8.2 mmol) heated to reflux in toluene (150 mL) for 3 hours. Yield 1.5 g (48.3 %). m.p. 250-253°C. ¹H NMR (DMSO-d₆) δ 12.75 (s, 1H), 7.98-7.97 (d, 1H, $J = 3.9$ Hz), 7.71-7.70 (d, 1H, $J = 5.2$ Hz), 7.65-7.62 (d, 1H, $J = 8.1$ Hz), 7.09-7.07 (d, 1H, $J = 8.1$ Hz), 2.13 2.08 (m, 6H), 1.99-1.85 (m, 2H); MS. [M-H]⁻ = 309. IR (SP ATR) 1430, 1620, 2220 cm⁻¹. Anal. C₁₇H₁₄N₂S₂. calc C, 65.77; H, 4.55; N, 9.02. obs'd C 65.27; H, 4.41; N, 8.84.

25

Example 13**5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione
5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-one:**

Prepared from 4-bromo-2-fluoroanisole and (2'-oxo-2',3'-dihydrospiro[cyclohexane-

30

- 69 -

1,3'-[3H]indol]-5'-yl) boronic acid according to the procedure for example 5 to afford the subtitled compound as a white solid, mp. 178 – 180 °C; ¹H-NMR (DMSO - d₆) δ 10.4 (s, 1H), 7.65 (d, 1H, *J* = 1.1 Hz), 7.5 - 7.4 (m, 3H), 7.2 (t, 1H, *J* = d *J* = 8.8 Hz), 3.9 (s, 3H), 1.9 (m, 2H) 1.7 - 1.6 (m, 8H); MS (APCI (-)) *m/z* 324 [M-H]⁺; Anal. Cal. For C₂₀H₂₀FNO₂ : C, 73.83, H, 6.20, N, 4.30. Found: C, 73.55, H, 6.23, N, 4.40.

The title compound was prepared by refluxing overnight a mixture of 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-one and an equal weight of phosphorus pentasulfide in pyridine. Removal of the pyridine *in vacuo* followed by treatment of the residue with 5N hydrochloric acid solution and subsequent recrystallization in ethanol gave a grey solid, mp 228-229°C; ¹H-NMR (DMSO-d₆) δ 12.7 (s, 1H), 7.9 (s, 1H), 7.6 - 7.5 (m, 2H), 7.5 - 7.4 (m, 1H), 7.2 (t, 1H, *J* = 8.8 Hz), 7.1 (d, 1H, *J* = 8.1 Hz), 3.9 (s, 3H), 1.9 - 1.8 (m, 7H), 1.4 - 1.3 (m, 3H); MS (APCI (-)) [M-H]⁺ *m/z* 324. Anal. Cal. for C₂₀H₂₀FNOS. 0.25 H₂O C, 69.44; H, 5.97; N, 4.05. Found: C, 69.43; H, 5.75; N, 4.32.

Example 14

5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H] indole]-2(1H)-thione

Prepared by refluxing overnight a mixture of 5-(2-amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]-indole]-2(1H)-one and an equal weight of phosphorus pentasulfide in pyridine. Removal of the pyridine *in vacuo* followed by treatment of the residue with 5N hydrochloric acid solution and subsequent recrystallization in ethanol gave a grey solid; mp 274 – 277 °C (dec.); ¹H-NMR (DMSO-d₆) δ 12.7 (s, 1H), 8.6 (s, 2H), 7.9 (s, 1H), 7.5 (d, 1H, *J* = 8.1 Hz), 7.1 (d, 1H, *J* = 8.1 Hz), 6.8 (s, 2H), 1.9 - 1.8 (m, 7H), 1.4 - 1.3 (m, 3H). MS (APCI (-)) [M-H]⁺ *m/z* 309.

Example 15**3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile****Spiro[cyclopentane-1,3'-[3H]indol]-2'(1'H)-one**

- 5 To a -25 °C solution of oxindole (2.0 g, 15.0 mmol) in 40 (cm³) of anhydrous THF under N₂ was added *n*-butyllithium (1.6 M in hexanes, 19.7 cm³, 31.5 mmol) drop-wise. To the resulting milky solution was added *N,N,N',N'*-tetramethylethylenediamine (4.75 cm³, 31.5 mmol). After 30 min. a solution of 1,4-diiodobutane (21.9 g, 70.6 mmol) in THF (3 cm³) was added and the reaction mixture
- 10 was allowed to warm to RT and stirred for 14 h. The reaction mixture was poured into water, extracted with EtOAc (x 2), the combined organic layers were washed with dilute HCl (pH 1) and water (x 2), dried (MgSO₄) and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc: hexane 1:4) to afford the subtitled compound (1.4 g, 7.5 mmol, 50%) as a tan solid: ¹H NMR (CDCl₃) δ 1.8-2.2 (m, 8H),
- 15 6.94 (dd, *J* = 7.5, 1.0 Hz, 1H), 7.01 (dd, *J* 7.5, 1.0 Hz, 1H), 7.14-7.25 (m, 2H), 9.30 (br s, 1H).

5-Bromo-spiro[cyclopentane-1,3'-[3H]indol]-2'(1'H)-one

- A solution of spiro[cyclopentane-1,3'-[3H]indol]-2'(1'H)-one (0.27 g, 1.4 mmol) and sodium acetate (0.12 g, 1.46 mmol) in acetic acid (10 cm³) was treated
- 20 with bromine (0.24 g, 1.51 mmol) in acetic acid (2 cm³). After 30 min. the mixture was poured into sat. sodium hydrogen carbonate solution and extracted with EtOAc (x 2), the combined organic layers were washed with water, sat. sodium hydrogen carbonate solution, water, dried (MgSO₄), and evaporated to give the subtitled compound (0.37 g, 1.47 mmol, 96 %) as an off-white solid which was used without
- 25 further purification: ¹H NMR (CDCl₃) δ 1.8-2.27 (m, 8H), 6.79 (d, *J* = 8 Hz, 1H), 7.30-7.39 (m, 2H), 8.63 (br s, 1H).

5'-(3-Cyano-5-fluorophenyl)-spiro[cyclopentane-1,3'-[3H]indol]-2'(1'H)-one:

- A solution of 3-cyano-5-fluoro-bromobenzene (0.5 g, 2.6 mmol), and tetrakis (triphenylphosphine) palladium(0) (0.2 g) in ethylene glycol dimethyl ether (20 cm³)
- 30 was stirred under N₂ for 20 minutes. To this mixture was then (spiro[cyclopentane-

- 71 -

1,3'-[3H]indol]-2'(1'H)-one-5-yl) boronic acid (0.9 g, 3.9 mmol) and sodium carbonate (0.8 g, 7.8 mmol) in water (5 cm³). The solution was brought to reflux for 18 hours and then cooled to room temperature, poured into 2N NaOH and extracted with EtOAc (x 3). The combined extracts were washed with water, brine, dried (MgSO₄), and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc, hexane) to afford the subtitled compound (0.35 g, 44%) as white needles. mp: 235 - 237 °C; ¹H NMR (DMSO-d₆) δ 10.5 (s, 1H), 8.1 (s, 1H), 8.0 (dt, 1H, *J* = 1.7, 2.0, 7.0 Hz), 7.8 - 7.7 (m, 2H), 7.6 (dd, 1H, *J* = 1.8, 6.4 Hz), 6.9 (d, 1H, *J* = 8.1 Hz), 2.0 - 1.9 (m, 8H); MS (EI) M⁺ @ m/z 306.

10 General Procedure A

The title compound was prepared from 5'-(3-Cyano-5-fluorophenyl)-spiro[cyclopentane-1,3'-[3H]indol]-2'(1'H)-one (40 mg) and Lawesson's reagent (50 mg) in toluene (10 ml) at reflux in a sealed tube for 16 h. The mixture was concentrated and the residue dissolved in a minimal amount of THF, then purified by HPLC (SiO₂, 30 cm x 2.5 cm, EtOAc-Hexane 2:8 at 20 ml/min.) to afford the title compound (0.022 g) as an off white solid: mp. 236 - 238 °C; ¹H NMR (DMSO-d₆) δ 12.66 (br s, 1H), 8.11 (s, 1H), 7.97 (dt, 1H, *J* = 10.1 and 2.2 Hz), 7.79 - 7.76 (m, 2H), 7.68 (dd, 1H, *J* = 8.1 and 1.7 Hz), 7.07 (d, 1H, *J* = 8.1 Hz), 2.10 - 2.05 (m, 6H) and 1.97 - 1.88 (m, 2H); MS (EI) m/z 322 [M]⁺.

20

Example 16

5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione

5-(3-Chloro-phenyl)-3,3-dimethyl-1,3-dihydro-indol-2-one. 5-bromo-1,3-dihydro-3,3-dimethyl-2H-indol-2-one (0.98 g, 4.07 mol) and tetrakis(triphenylphosphine)palladium(0) (0.239 g) were stirred under an atmosphere of nitrogen in dimethoxyethane (35 cm³). After 15 min., 3-chlorophenylboronic acid (1.27 g, 8.13 mol) was added, followed by potassium carbonate (3.40 g, 45 mmol) in water (15 cm³). The reaction was heated to reflux for 2 hours and then stirred at room temperature overnight. The mixture was diluted with sat. ammonium chloride and extracted with EtOAc (x3). The combined organic layers were dried (MgSO₄),

30

- 72 -

filtered, and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc: hexane, 1:3) to afford the subtitled compound (0.284 g, 25%): mp 188 -189 °C; ¹H NMR (DMSO-d₆) δ 3.34 (s, 6 H), 6.93 (d, 1 H, *J* = 8.04 Hz), 7.38-7.35 (m, 1 H), 7.53-7.43 (m, 2 H), 7.61 (d, 1 H, *J* = 7.68 Hz), 7.70 (s, 2 H), 10.40 (s, 1 H); IR (KBr) 3420, 3150, 3050, 1700 cm⁻¹; MS (EI) *m/z* 270 (M-H)⁺; CHN calculated for C₁₆H₁₄ClNO + 0.1C₄H₈O₂: C, 70.21; H, 5.32; N, 4.99; Found: C, 70.3; H, 5.44; N, 4.93.

The title compound was prepared from 5-(3-Chloro-phenyl)-3,3-dimethyl-1,3-dihydro-indol-2-one (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.031 g) as an off white solid: mp. 158 – 160 °C; ¹H NMR (CDCl₃) δ 9.67 (br s, 1H), 7.55 (s, 1H), 7.47 – 7.43 (m, 3H), 7.40 – 7.30 (m, 2H), 7.08 (d, 1H, *J* = 8.7 Hz) and 1.50 (s, 6H); MS (EI) *m/z* 287/289 [M]⁺.

15

Example 17

3-Benzyl-5-(3-chlorophenyl)-3-methyl-1,3-dihydro-2H-indole-2-thione

The title compound was prepared from 3-benzyl-5-(3-chloro-phenyl)-3-methyl-1,3-dihydro-indol-2-one (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.022 g) as an off white solid: mp. 168 – 170 °C; ¹H NMR (CDCl₃) δ 9.23 (br s, 1H), 7.49 (s, 1H), 7.49 – 7.30 (m, 4H), 7.21 (s, 1H), 7.15 – 7.09 (m, 3H), 6.96 – 6.94 (m, 2H), 6.89 (d, 1H, *J* = 8.0 Hz), 3.19 (dd, 2H, *J* = 40.5 and 13 Hz) and 1.57 (s, 3H); MS (EI) *m/z* 363/365 [M]⁺.

25

Example 18

4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile 4-(3,3-Dimethyl-2-oxo-2,3-dihydro-1H-indol-5-yl)-furan-2-carbonitrile.

Prepared according to the procedure for Example 5 using (2'-oxo-[2, 3-dihydro-3,3-dimethyl -1, 3'- [3H] indol] -5'-yl) boronic acid (354 mg, 1.7 mmol) and 4-bromo-furan-2-carbonitrile (200 mg, 1.2 mmol) to afford the subtitled compound (76 mg, 0.3

- 73 -

mmol, 26 %) as a white solid: mp. 199.6-201.4 °C, ¹H NMR (DMSO-d₆) δ 1.28 (s, 6H), 6.89 (d, *J* = 8.0 Hz, 1H), 7.48 (dd, *J* = 8.0, 1.8 Hz, 1H), 7.65 (d, *J* = 1.5 Hz, 1H), 8.1 (s, 1H), 8.5 (s, 1H), 10.46 (s, 1H); MS (ESI) *m/z* 251 (M-H)⁺; Anal. C₁₅H₁₂N₂O₂·0.6 H₂O

5 The title compound was prepared from 4-(3,3-Dimethyl-2-oxo-2,3-dihydro-1H-indol-5-yl)-furan-2-carbonitrile (73 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.003 g) as an off white solid: mp. 188 – 191 °C; ¹H NMR (CDCl₃) δ 9.63 (br s, 1H), 7.83 (s, 1H), 7.36 – 7.33 (m, 3H), 7.06 (d, 1H, *J* = 7.9 Hz) and 1.48 (s, 6H); MS (EI) *m/z* 268 [M]⁺.

10

Example 19

5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione

5-bromo-1,3-dihydro-3,3-dimethyl-2H-indol-2-one: 3,3-dimethyl-indol-2-one (0.65 g, 4.03 mmol) and sodium acetate (0.33 g, 4.07 mmol) were stirred in acetic acid (5 cm³) then bromine (0.66 g, 4.13 mmol) in acetic acid (5 cm³) was added drop-wise to the reaction mixture. The reaction was stirred for 50 min., then poured into water. The mixture was basified with sodium carbonate, extracted with ethyl acetate (x3), dried (MgSO₄), filtered, and evaporated to give the subtitled compound (0.89 g, 92%) ¹H NMR (DMSO-d₆) δ 1.21 (s, 6 H), 6.76 (d, 1 H, *J* = 8.22 Hz), 7.29 (dd, 1 H, *J* = 2.12 Hz, 8.23 Hz), 7.49 (d, 1 H, *J* = 2.03 Hz), 10.4 (s, 1H).

15

20

5-bromo-1,3-dihydro-3,3-dimethyl-2H-indol-2-one (0.33 g, 1.38 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.094 g) were stirred under an atmosphere of nitrogen in dimethoxyethane (12 cm³). After 15 minutes, 3-methoxyphenylboronic acid (0.42 g, 2.76 mmol) was added, followed by potassium carbonate (1.15 g, 8.34 mmol) in water (5 cm³). The reaction was heated to reflux for 5 hours, and then cooled to room temperature. Saturated aqueous ammonium chloride and EtOAc were added and the mixture was filtered. The aqueous layer was extracted with EtOAc (x2), and the combined organic layers were dried (MgSO₄), filtered, and evaporated. The residue was purified by column

25

30

- 74 -

chromatography (SiO₂, EtOAc: hexane 1:3) to afford 5-(3-methoxy-phenyl)-3,3-dimethyl-1,3-dihydro-indol-2-one (0.11 g, 31%), mp = 157-158 °C; ¹H NMR (DMSO-d₆) δ 3.34 (s, 6 H), 3.82 (s, 3 H), 6.87 - 6.93 (m, 2 H), 7.20-7.15 (m, 2 H), 7.37-7.32 (m, 1 H), 7.49-7.46 (m, 1 H), 7.63 (d, 1 H, *J* = 1.14 Hz), 10.4 (s, 1 H); MS (EI) *m/z* 266 (M-H)⁺; CHN calculated for C₁₇H₁₇NO₂: C, 76.38; H, 6.41; N, 5.24; Found: C, 76.02; H, 6.49; N, 5.02.

The title compound was prepared from 5-(3-methoxy-phenyl)-3,3-dimethyl-1,3-dihydro-indol-2-one (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.022 g) as an off white solid: mp. 149 - 150 °C; ¹H NMR (CDCl₃) δ 9.69 (br s, 1H), 7.49 - 7.46 (m, 2H), 7.37 (t, 1H, *J* = 8.0 Hz), 7.16 (d, 1H, *J* = 7.7 Hz), 7.09- 7.06 (m, 2H), 6.90 (dd, 1H, *J* = 8.2 and 2.3 Hz) 3.88 (s, 3H) and 1.50 (s, 6H); MS (EI) *m/z* 283 [M]⁺.

15

Example 20**3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile****3-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile**

Prepared according to the procedure for Example 5: m.p. 205 - 206 °C. ¹H NMR (DMSO-d₆) δ 10.47 (s, 1H), 8.08 - 8.06 (dd, 1H), 7.89 - 7.85 (m, 1H), 7.65 (s, 1H), 7.54 - 7.49 (m, 1H), 7.43 - 7.40 (tt, 1H), 6.95 - 6.93 (d, 1H *J* = 7.9 Hz), 1.97 - 1.83 (m, 2H), 1.69 - 1.55 (m, 8H); MS (EI) *m/z* 320 (M⁺).

The title compound was prepared from 3-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.037 g) as an off white solid: mp. 230 - 233 °C; ¹H NMR (CDCl₃) δ 9.82 (br s, 1H), 7.86 (s, 1H), 7.77 (dd, 1H, *J* = 7.0 and 1.8 Hz), 7.68 - 7.63 (m, 1H), 7.45 (d, 1H, *J* = 8.0 Hz), 7.31 (d, 1H, *J* = 9.0 Hz), 7.15 (d, 1H, *J* = 8.1 Hz), 2.17 - 1.84 (m, 7 H) and 1.60 - 1.54 (m, 3H); MS (EI) *m/z* 336 [M]⁺.

30

Example 21**5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile**

A solution of 3-bromopyridine-5-carbonitrile (2.79 g, 15.26 mmol),
5 hexamethylditin (5.00 g, 15.26 mmol) and tetrakis(triphenylphosphine)palladium(0)
(0.20 g, 0.17 mmol) in anhydrous dimethoxyethane (30 cm³) under N₂ was heated
under reflux. After 16 h the mixture was concentrated and purified by column
chromatography (SiO₂, EtOAc: hexane 5:95) to afford 3-cyanopyridine-5-
trimethylstannane (2.82 g, 10.55 mmol, 69%): ¹H NMR (CDCl₃) δ 0.40 (s, 9H), 8.01
10 (m, 1H), 8.80 (m, 2H); MS ((+) APCI) m/z 269 (M + H)⁺.

A solution of 5'-bromospiro[cyclohexane-1,3'-[3H] indol]-2'-(1'H)-one (1.97
g, 7.05 mmol), 3-cyanopyridine-5-trimethylstannane (2.26 g, 8.46 mmol),
bis(triphenylphosphine)palladium(II)chloride (0.33 g, 0.47 mmol) and lithium
chloride (1.48 g, 35 mmol) in anhydrous toluene (30 cm³) was heated under reflux.
15 After 16h the mixture was cooled, partitioned between EtOAc and water, the aqueous
layer was re-extracted with EtOAc (x 2), the combined organic extracts were washed
with water, dried (MgSO₄) and evaporated. The residue was subjected to column
chromatography (SiO₂, EtOAc: hexane, 1:2) and then further purified by preparative
LC (Primesphere C18, 10 micron, 50 x 250 mm, MeCN: H₂O 1:1, 100 cm³/min., RT
20 7.92 min.) to afford 3-(1',2'-dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol-5'-
yl)pyridine carbonitrile. as white crystals (0.56 g, 1.84 mmol, 26%): mp. 232 - 234
°C, ¹H NMR (CDCl₃) δ 1.68 - 1.89 (m, 6H), 1.93 - 2.13 (m, 4H), 7.12 (d, 1H, J = 8
Hz), 7.49 (dd, 1H, J = 8, 2 Hz), 7.66 (d, 1H, 2 Hz), 8.15 (t, 1H, J = 2 Hz), 8.39 (s, 1H,
br), 8.89 (d, 1H, J = 2 Hz), 9.06 (d, 1H, J = 2 Hz); MS ((+)-ESI) m/z 304 (M + H)⁺;
25 Anal. C₁₉H₁₇N₃O CHN.

The title compound was prepared from 3-(1',2'-Dihydro-2'-
oxospiro[cyclohexane-1,3'-[3H]indol-5'-yl)pyridine carbonitrile (100 mg) and
Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General
Procedure A, to afford the title compound (0.004 g) as a yellow solid: mp. 237 - 238
30 °C; ¹H NMR (CDCl₃) δ 9.56 (br s, 1H), 9.03 (d, 1H, J = 1.9 Hz), 8.87 (d, 1H, J = 1.4

- 76 -

Hz), 8.12 (s, 1H), 7.87 (s, 1H), 7.50 (d, 1H, $J = 8.1$ Hz), 7.17 (d, 1H, $J = 8.1$ Hz), 2.19 – 1.85 (m, 7H) and 1.59 – 1.54 (m, 3H); MS ((-)-APCI) m/z 318 [M-H]⁻.

Example 22

5 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione
 5'-(3,5-Difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one:

Prepared according to the procedure for Example 5: mp 180-183 °C; ¹H-NMR (CDCl₃) δ 8.35 (s, 1H), 7.59 (d, 1H, $J = 2.0$ Hz), 7.40 (dd, 1H, $J = 6.2, 2.0$ Hz), 7.10 - 7.03 (m, 2H), 6.99 (d, 1H, $J = 8.1$ Hz), 7.76 (tt, 1H, $J = 4.3, 2.3$ Hz), 2.05 - 1.62 (m, 10H); MS ((+)-APCI) m/z 314 [M+H]⁺.

10 The title compound was prepared from 5'-(3,5-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.020 g) as a yellow solid: mp. 232 – 233 °C; ¹H NMR (CDCl₃) δ 10.05 (br s, 1H), 7.83 (s, 1H), 7.44 (dd, 1H, $J = 8.1$ and 1.4 Hz), 7.38 – 7.30 (m, 1H), 7.26 – 7.19 (m, 3H), 7.11 (d, 1H, $J = 8.1$ Hz), 2.17 – 1.82 (m, 7H) and 1.66 – 1.53 (m, 3H); MS ((-)-APCI) m/z 328 [M-H]⁻.

Example 23

20 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione
 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H] indol]-2(1H)-one:

Prepared according to the procedure for Example 5: m.p. 191-192°C, ¹H NMR (CDCl₃) δ 1.6-2.1 (m, 10H), 6.85-6.95 (m, 2H), 6.98 (d, $J = 4.0$ Hz, 1H), 7.36 (dd, $J = 7.5, 1.6$ Hz, 1H), 7.53 (d, $J = 0.9$ Hz, 1H), 7.80 (br s, 1H); ¹³C-NMR (THF-d₈) δ 21.35, 25.33, 33.12 (t), 48.32 (s), 110.40, 121.66, 121.96, 125.44, 127.25 (d), 128.17, 128.43, 136.92, 140.20, 143.43, 183.72 (s); MS (EI) m/z 318 (M+H)⁺; Anal. (C₁₇H₁₆ClNOS) C, H, N.

25 The title compound was prepared from 5-(5-Chloro-2-thienyl)spiro [cyclohexane-1,3-[3H] indol]-2(1H)-one (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product

30

- 77 -

(0.041 g) as a yellow solid: mp. 231 – 232 °C; ¹H NMR (CDCl₃) δ 9.75 (br s, 1H), 7.82 (d, 1H, *J* = 1.2 Hz), 7.43 (dd, 1H, *J* = 8.1 and 1.6 Hz), 7.04 – 7.02 (m, 2H), 6.89 (d, 1H, *J* = 3.8), 2.15 – 1.84 (m, 7H) and 1.59 – 1.52 (m, 3H); MS ((-)-APCI) *m/z* 332/334 [M-H]⁺.

5

Example 24

5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile

5-(1',2'-Dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-3-

10 **furancarbonitrile:**

Prepared according to the procedure for Example 5: m.p. 243 – 245 °C. ¹H-NMR (DMSO-d₆) δ 10.48 (s, 1H), 8.62 (d, 1H *J* = 0.7 Hz), 7.76 (d, 1H *J* = 1.5 Hz), 7.58 - 7.55 (dd, 1H), 7.33 (d, 1H *J* = 0.7 Hz), 6.92 - 6.90 (d, 1H *J* = 8.1 Hz), 1.87 - 1.83 (m, 2H), 1.73 - 1.53 (m, 8H). MS ((+)EI) *m/z* 292 (M⁺).

15

The title compound was prepared from 5-(1',2'-dihydro-2'-oxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-3-furancarbonitrile (100 mg) and Lawesson's reagent (120 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.020 g) as a yellow solid: mp. 264 – 268 °C; ¹H NMR (CDCl₃) δ 9.66 (br s, 1H), 7.98 (s, 2H), 7.59 (dd, 1H, *J* = 8.2 and 1.5 Hz), 7.08 (d, 1H, *J* = 8.2 Hz), 6.78 (s, 1H), 2.16 – 1.85 (m, 7H) and 1.56 – 1.52 (m, 2H); MS ((-)-APCI) *m/z* 307 [M-H]⁺.

20

Example 25

5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione

25 **5'-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one.**

Prepared according to the procedure for Example 5: mp 188-189 °C; ¹H-NMR (CDCl₃) δ 7.97 (s, 1H), 7.57 - 7.54 (m, 2H), 7.41 - 7.34 (m, 2H), 7.20 (t, 1H, *J* = 8.7 Hz), 9.96 (d, 1H, *J* = 8.1 Hz), 2.04 - 1.65 (m, 10H); MS ((+)APCI) *m/z* 330 [M+H]⁺.

The title compound was prepared from 5'-(3-Chloro-4-fluorophenyl)

- 78 -

spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (100 mg) and Lawesson's reagent (100 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.036 g) as an off white solid: ¹H NMR (DMSO-d₆) δ 12.74 (br s, 1H), 7.92 (d, 1H, *J* = 1.4 Hz), 7.87 (dd, 1H, *J* = 7.1 and 2.3 Hz), 7.70 – 7.65 (m, 1H), 7.61 (dd, 1H, *J* = 7.1 and 1.5 Hz), 7.49 (t, 1H, *J* = 8.9 Hz), 7.14 (d, 1H, *J* = 8.1 Hz), 1.99 – 1.82 (m, 7H) and 1.40 – 1.37 (m, 3H); MS ((-)-APCI) *m/z* 344/346 [M-H]⁻.

Example 26

5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione

10 5'-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one:

Prepared according to the procedure for Example 5: mp 178-180 °C; ¹H-NMR (CDCl₃) δ 8.50 (s, 1H), 7.57 (d, 1H, *J* = 1.8 Hz), 7.39 (dd, 1H, *J* = 6.2, 1.9 Hz), 7.33 - 7.32 (m, 1 H), 7.15 (dq, 1H, *J* = 5.7, 1.7, 0.7 Hz), 7.06 (dq, 1 H, *J* = 4.2, 1.9, 0.4 Hz), 7.00 (d, 1H, *J* = 8.1 Hz), 2.05 - 1.64 (m, 10H); MS ((-)-ESI) [M-H]⁻ @ *m/z* 328.

15 The title compound was prepared from 5'-(3-chloro-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (100 mg) and Lawesson's reagent (100 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.039 g) as an off white solid: ¹H NMR (DMSO-d₆) δ 12.76 (br s, 1H), 7.97 (d, 1H, *J* = 1.1 Hz), 7.67 (dd, 1H, *J* = 8.1 and 1.4 Hz), 7.60 – 7.54 (m, 2H), 7.40 (dt, 1H, *J* = 8.65 and 2.0 Hz), 7.14 (d, 1H, *J* = 8.1 Hz), 1.99 – 1.83 (m, 7H) and 1.41 – 1.38 (m, 3H); MS ((-)-APCI) *m/z* 344/346 [M-H]⁻.

Example 27

5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione

25 5'-(3,5-Difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one:

Prepared according to the procedure for Example 5: mp 180-183 °C; ¹H-NMR (CDCl₃) δ 8.35 (s, 1H), 7.59 (d, 1H, *J* = 2.0 Hz), 7.40 (dd, 1H, *J* = 6.2, 2.0 Hz), 7.10 - 7.03 (m, 2H), 6.99 (d, 1H, *J* = 8.1 Hz), 7.76 (tt, 1 H, *J* = 4.3, 2.3 Hz), 2.05 - 1.62 (m, 10H); MS ((+)-APCI) *m/z* 314 [M+H]⁺.

- 79 -

The title compound was prepared from 5'-(3,5-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (100 mg) and Lawesson's reagent (100 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound 0.029 g as an off white solid: ¹H NMR (DMSO-d₆) δ 12.76 (br s, 1H), 7.84 (s, 1H), 7.64 – 7.56 (m, 1H), 7.46 (d, 1H, *J* = 8.1 Hz), 7.40 – 7.32 (m, 1H), 7.22 – 7.15 (m, 2H), 1.99 – 1.80 (m, 7H) and 1.38 – 1.35 (m, 3H); MS ((-)-APCI) *m/z* 328 [M-H]⁺.

Example 28

10 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile

5-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile. The title compound was prepared in a manner similar to Example 5 from 5-bromo-4-n-propyl thiophene-2-carbonitrile (1.17 g, 5 mmol), (1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H] indol]-5-boronic acid (1.24 g, 5 mmol), tetrakis(triphenylphosphine) palladium, potassium carbonate (2.75 g, 21 mmol), water (10 mL), and dimethoxyethane (50 mL) heated at reflux over night, to afford the product (0.7 g, 40%): m.p. 168-171 °C; ¹H NMR (DMSO-d₆) δ 10.56 (s, 1H), 7.93 (s, 1H) 7.52-7.51 (d, 1H, *J* = 1.5 Hz), 7.33 - 7.29 (dd, 1H, *J* = 1.6 Hz), 7.00-6.96 (d, 1H, *J* = 8.0 Hz), 2.62-2.57 (t, 2H), 1.86 (m, 2H), 1.70-1.56 (m, 11 H), 0.88-0.84 (t, H); MS *m/z* (APCI (+)) 351 [M+H]⁺. IR (KBr) 1620, 1700, 2200 cm⁻¹. Anal. C₂₁H₂₂N₂OS·1/2 H₂O calc. C, 70.2; H, 6.39; N, 7.79. Observed. C, 70.67; H, 6.34; N, 7.62.

The title compound was prepared from 5-(1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl- 2-thiophenecarbonitrile (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.037 g) as an orange solid: ¹H NMR (DMSO-d₆) δ 12.83 (br s, 1H), 7.96 (s, 1H), 7.77 (s, 1H), 7.44 (d, 1H, *J* = 7.7 Hz), 7.19 (d, 1H, *J* = 8.0 Hz), 2.60 (t, 2H, *J* = 8.0 Hz), 1.98 – 1.79 (m, 7H), 1.64 – 1.56 (m, 2H), 1.39 – 1.35 (m, 2H) and 0.87 (t, 3H, *J* = 7.3 Hz): MS ((-)-APCI) *m/z* 365 [M-H]⁺.

Example 29**5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione****5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-one:**

- 5 Prepared from (2'-oxo-2,3-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)boronic acid (3.2 g, 12.5 mmol) and 4-bromo-2-fluoro-nitrobenzene (3 g, 13.6 mmol) as described for example 5, to afford the title compound (0.7 g, 16%) as a yellow solid: mp. 213-215 °C; ¹H NMR (DMSO-d₆) δ 1.5 - 1.8 (m, 8H), 1.8 - 2.0 (m, 2H), 6.96 (d, 1H, *J* = 8.13 Hz), 7.68 (dd, 1H, *J* = 8.13, 1.76 Hz), 7.74 (dd, 1 H, *J* = 8.68, 1.76 Hz),
10 7.86 (d, 1H, *J* = 1.98 Hz), 7.92 (dd, 1H, *J* = 13.4, 1.76 Hz), 8.18 (t, 1H, *J* = 8.46 Hz) and 10.52 (s, 1H); MS (EI) *m/z* = 340 (M⁺).

- The title compound was prepared from 5-(3-fluoro-4-nitrophenyl)spiro [cyclohexane-1,3-[3H]indol]-2(1H)-one (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product
15 (0.021 g) as a yellow solid: ¹H NMR (DMSO-d₆) δ 12.82 (br s, 1H), 8.21 (t, 1H, *J* = 8.4 Hz), 8.07 (d, 1H, *J* = 1 Hz), 7.98 (dd, 1H, *J* = 13.1 Hz), 7.79 (dt, 1H, *J* = 8.1 and 2.6 Hz), 7.19 (1H, *J* = 8.2 Hz), 1.99 - 1.83 (m, 7H) and 1.42 - 1.39 (m, 3H); MS ((-)-APCI) *m/z* 355 [M-H].

20

Example 30**4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile**

- 4-(1,2-Dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile: A solution of 3-bromo-5-cyano-furan (0.75 g, 4.4 mmol), and
25 tetrakis(triphenylphosphine) palladium(0) (0.4 g) in ethylene glycol dimethyl ether (20 cm³) was stirred under N₂ for 20 minutes. To this mixture was then added (spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one-5-yl) boronic acid (1.6 g, 6.5 mmol) and sodium acetate (1.4 g, 13.1 mmol) in water (5 cm³). The solution was brought to reflux for 18 hours and then cooled to room temperature, poured into 2N NaOH and
30 extracted with EtOAc (x 3). The combined extracts were washed with water, brine,

- 81 -

dried (MgSO₄), and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc, hexane) to afford the product (0.45 g, 36%) as an off-white solid. mp: 240 - 242 °C; ¹H NMR (DMSO-d₆) δ 10.4 (s, 1H), 8.5 (s, 1H), 8.2 (s, 1H), 7.7 (s, 1H), 7.5 (dd, 1H, *J* = 1.5 6.5 Hz), 6.9 (d, 1H, *J* = 8.0 Hz), 2.0 - 1.6 (m, 10H); MS (EI) M⁺ @ *m/z* 292.

The title compound was prepared from 4-(1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile (67 mg) and Lawesson's reagent (67 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.018 g) as a yellow solid: ¹H NMR (DMSO-d₆) δ 12.74 (s, 1H), 8.68 (s, 1H), 8.26 (s, 1H), 7.96 (s, 1H), 7.62 (dd, 1H, *J* = 8.0 and 1.0 Hz), 7.10 (s, 1H, *J* = 8.1 Hz), 1.94 - 1.78 (m, 7H) and 1.35 - 1.32 (m, 3H); MS ((-)-APCI) *m/z* 307 [M-H].

Example 31

15 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indole]-2''(1''H)-thione

5-Bromospiro[cyclobutane-1,3-[3H]indol]-2(1H)-one: To a stirred solution of spiro[cyclobutane-1,3'-[3H]indol]-2'(1'H)-one (*J. Med. Chem.* **1987**, 824-9) (1.0 g, 6 mmol) in glacial acetic acid (10 mL) was added dropwise at room temperature a solution of bromine (0.30 mL, 6 mmol) in glacial acetic acid (6 mL). After stirring 20 for 10 min, anhydrous sodium acetate (0.47 g, 6 mmol) was added and the solution was concentrated *in vacuo*. The residue was dissolved in ethyl ether (50 mL) and washed sequentially with water (50 mL), aqueous saturated sodium bicarbonate solution (50 mL), water (50 mL) and brine (30 mL). The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. Crystallization from ethyl 25 ether yielded the product as a white fluffy solid (1.1 g, 73%), mp 235-7 °C. ¹H NMR (DMSO-d₆, 300 MHz) 2.15-2.41 (m, 6 H), 6.74 (d, 1 H, *J* = 8.2 Hz), 7.33 (dd, 1 H, *J* = 2, 8.2 Hz), 7.75 (d, 1 H, *J* = 2 Hz), 10.36 (bs, 1 H). MS (EI) *m/z* 251 [M⁺]. Anal. Calcd for C₁₁H₁₀BrNO: C, 52.41; H, 4.00; N, 5.56. Found: C, 51.98; H, 4.24; N, 5.42.

- 82 -

To a solution of 5-bromospiro[cyclobutane-1,3-[3H]indol]-2(1H)-one (0.6 g, 2 mmol) in ethylene glycol dimethyl ether (50 mL) under a nitrogen atmosphere was added tetrakis(triphenylphosphine)palladium(0) (140 mg, 0.1 mmol). To the solution was added sequentially 3-chlorophenyl boronic acid (0.48 g, 3 mmol) and potassium carbonate (0.76 g, 5 mmol) in water (5 mL). The mixture was heated to 80 °C for 3 h and allowed to cool. The reaction mixture was poured into water (100 mL) and extracted with ethyl acetate (3 x 100 mL). The organic layers were combined, washed with brine (50 mL) and dried over magnesium sulfate. The solution was filtered, concentrated *in vacuo*, and the residue was purified by HPLC (Zorbax PRO, C18, 10u, 15A, 50 X 250mm; 35% Water/65% AcCN; 254NM; AMB. temp.) to give 5-(3-chlorophenyl)spiro[cyclobutane-1,3-[3H]indole]-2(1H)-one (200 mg, 35%) as a white powder, mp 199.5-201 °C. ¹H NMR (DMSO-*d*₆, 300 MHz) 2.21-2.28m, 2 H), 2.40-2.45 (m, 4 H), 6.87 (d, 1 H, *J* = 8.1 Hz), 7.37 ('d', 1 H), 7.44-7.52 (m, 2 H), 7.65 (bd, 1 H, *J* = 7.8 Hz), 7.76 (bs, 1 H), 7.92 (bs, 1 H), 10.35 (s, 1 H). MS (EI) *m/z* 283 [M⁺]. Anal. Calcd for C₁₇H₁₄ClNO: C, 71.96; H, 4.97; N, 4.94. Found: C, 70.75; H, 5.07; N, 4.68.

The title compound was prepared from 5-(3-Chlorophenyl)spiro[cyclobutane-1,3-[3H]indole]-2(1H)-one (55 mg) and Lawesson's reagent (55 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound 0.016 g as an orange solid: ¹H NMR (DMSO-*d*₆) δ 12.58 (br s, 1H), 8.07 (d, 1H, *J* = 1.5 Hz), 7.82 (t, 1H, *J* = 1.7 Hz), 7.70 (d, 1H, *J* = 7.74 Hz), 7.60 (dd, 1H, *J* = 8.12 and 1.71 Hz), 7.49 (t, 1H, 7.9 Hz), 7.41 (d, 1H, *J* = 8.32 Hz), 7.05 (d, 1H, *J* = 8.14 Hz) and 2.57 – 2.27 (m, 6H); MS ((-)-APCI) *m/z* 298/300 [M-H].

25

Example 32**5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indole]-2''(1''H)-thione**

The title compound was prepared from 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indole]-2''(1''H)-thione (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product 0.042 g as an off white solid: ¹H NMR (DMSO-*d*₆)

30

- 83 -

δ 12.75 (br s, 1H), 7.80 (d, 1H, $J = 1.1$ Hz) 7.58 – 7.55 (m, 1H), 7.48 – 7.36 (m, 4H), 7.16 (d, 1H, $J = 8.0$ Hz) ; MS ((-)-APCI) m/z 326/328 [M-H]⁻.

Example 33

5 **5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indole-2''(1''H)-thione**

The title compound was prepared from 5-(4-chlorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-one (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product 0.035 g as an off white solid: ¹H NMR (DMSO-d₆) δ 12.74 (br s, 1H), 7.91 (d, 1H, $J = 1.3$ Hz), 7.69 (d, 2H, $J = 5.5$ Hz), 7.60 (dd, 1H, $J = 8.1$ and 1.4 Hz), 7.50 (d, 2H, $J = 8.5$ Hz), 10 7.15 (d, 1H, $J = 8.1$ Hz), 1.99 – 1.83 (m, 7H) and 1.50 – 1.36 (m, 3H); MS ((-)-APCI) m/z 326/328 [M-H]⁻.

Example 34

15 **5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)- 4-methyl-2-thiophenecarbonitrile**

5-Bromo-4-methyl-2-thiophene carboxaldehyde: To a solution of diethylamine (28g, 0.383 mol) in anhydrous THF (400 mL) was added at -40 °C under nitrogen a solution of *n*-BuLi (2.5 M, 153 mL, 0.383 mol) in hexane. After addition, the solution was stirred at -40 °C under nitrogen for 30 minutes, cooled to - 20 78 °C and treated dropwise with a solution of 2-bromo-3-methylthiophene (45g, 0.254 mol) in anhydrous THF (450 mL). The reaction solution was stirred at -78 °C for 30 minutes and treated with anhydrous DMF (100 mL). The mixture was allowed to warm to ambient temperature and was quenched with 1N aqueous hydrochloride solution (1L). The solution was extracted with ethyl acetate (3x450 mL) and the 25 extracts washed with water, brine and dried (MgSO₄). After removal of solvent *in vacuo*, the title compound was obtained as a white solid (46g, 88.3%). A sample of the product was crystallized from hexane: mp 63-65 °C; IR (KBr) 1654 cm⁻¹. ¹H-NMR (CDCl₃) δ 9.75 (s, 1H), 7.45 (s, 1H), 2.26 (s, 3H); MS (EI) m/z 204/206 (M⁺). 30 Anal. Calc. For C₆H₇BrOS: C, 35.14; H, 2.46. Found: C, 35.00; H, 2.44.

5-Bromo-4-methyl-2-thiophenecarbonitrile: Prepared from 5-bromo-4-methyl-2-thiophene carboxaldehyde using the procedure of Example 5. White solid: mp 40-42 °C; IR (KBr) 2200 cm⁻¹; ¹H-NMR (CDCl₃) δ 7.29 (s, 1H), 2.21 (s, 3H). MS (EI) *m/z* 201/203 (M⁺, 98%/100%); Anal. Calc. For C₆H₄BrNS: C, 35.66; H, 1.99; N, 6.93. Found: C, 36.00; H, 2.14; N, 6.76.

Prepared according to the procedure for Example 5 using (2'-oxo-[2, 3-dihydro-3,3-dimethyl -1, 3'-[3H] indol] -5'-yl) boronic acid (357 mg, 1.7 mmol) and 5-bromo-4-methylthiophene-2-carbonitrile (295 mg, 1.5 mmol) to afford 5-(3,3-Dimethyl-2-oxo-2,3-dihydro-1H-indol-5-yl)-4-methyl thiophene-2-carbonitrile (227 mg, 0.8 mmol, 55 %) as a white solid: mp. 192.3-193 °C, ¹H NMR (DMSO-d₆) δ 1.29 (s, 6H), 2.29 (s, 3H), 6.97 (d, *J* = 8.0 Hz, 1H), 7.34 (dd, *J* = 8.0, 1.8 Hz, 1H), 7.49 (d, *J* = 1.7 Hz, 1H), 7.84 (s, 1H), 10.57 (s, 1H); MS (EI) *m/z* 282 (m)⁺; Anal. C₁₆H₁₄N₂OS.

The title compound was prepared from 5-(3,3-dimethyl-2-oxo-2,3-dihydro-1H-indol-5-yl)-4-methyl thiophene-2-carbonitrile (0.77 g, 2.39 mmol) and phosphorous pentasulfide (0.42 g, 0.96 mmol) in toluene (20 ml) at reflux. After 3 h, the reaction was cooled and partitioned between water and EtOAc, the organic layer was separated, dried (MgSO₄) and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc- hexane gradient elution) to afford the product (0.25 g, 0.73 mmol, 30%) as an orange solid: ¹H NMR (DMSO-d₆) δ 12.82 (br s, 1H), 7.88 (s, 1H), 7.82 (d, 1H, 2 Hz), 7.49 (dd, 1H, *J* = 8.1, 1.6 Hz), 7.18 (d, 1H, *J* = 8.1 Hz), 1.99 – 1.80 (m, 7H) and 1.40 – 1.36 (m, 3H); MS ((-)-APCI) *m/z* 321 [M-H]⁻.

Example 35

5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)- 2-thiophenecarbonitrile

5-Bromo-2-thiophenecarbonitrile: A mixture of 5-bromo-2-thiophenecarboxaldehyde (96.0g, 500 mmol), hydroxylamine hydrochloride (111.9 g, 500 mmol), pyridine (500 mL), and ethanol (500 mL) was heated under nitrogen at reflux for two hours. The reaction mixture was cooled to ambient temperature and

concentrated *in vacuo* to give an oil. The crude product was triturated twice with ice water and the solid obtained was collected on a filter. A mixture of a portion of the above solid (44.31 g, 215 mmol), copper (II) acetate monohydrate (4.2 g, 21 mmol) in acetonitrile (1.4L) was heated at reflux for three hours. The solvent was removed *in vacuo* and the residue was dissolved in ethyl acetate. The solution was washed with 5% aqueous sulfuric acid (2X30 mL), water (2X30 mL), brine (20 mL), and dried (MgSO₄). The solvent was removed *in vacuo* and the residue was dissolved in a minimum amount of chloroform (1L) and allowed to crystallize. The crystals obtained was collected on a filter and the filtrate was concentrated and purified by a chromatography (silica gel, chloroform) to give the subtitled compound as an off-white solid (31.5g combined, 58%). IR (film) cm⁻¹ 2200. ¹H-NMR (CDCl₃) δ 7.39-7.38 (d, 1H, *J* = 4.1 Hz), 7.10 (d, 1H, *J* = 4.0 Hz); MS (EI) *m/z* 187 (M⁺, 98%) 189(M⁺, 100%).

5-(2'-Oxo-2',3'-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile was prepared according to the procedure for Example 5 using 5-bromo-2-thiophenecarbonitrile and (2'-oxo-2',3'-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl) boronic acid: mp. 225-228°C; ¹H NMR (DMSO-d₆) δ 1.63 (m, 8H), 1.90 (m, 2H) 6.91 (d, 1H, *J* = 8.13 Hz), 7.55 (dd, 1H, *J* = 8.13, 1.76Hz), 7.60 (d, 1H, *J* = 4.17 Hz), 7.75 (d, 1H, *J* = 1.76 Hz), 7.93 (d, 1H, *J* = 4.17 Hz), 10.51 (s, 1H); MS ((+)APCI) *m/z* 309 [M + H]⁺.

The title compound was prepared from 5-(2'-oxo-2',3'-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile (0.69 g) and phosphorous pentasulfide (0.4 g) in toluene (20 ml) at reflux. After 3 h. the reaction was cooled, poured into sat. aqueous sodium hydrogen carbonate solution, and extracted with EtOAc. The organic layer was separated, dried (MgSO₄) and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc-hexane gradient elution) to afford the title compound (0.215 g) as an orange solid: ¹H NMR (DMSO-d₆) δ 12.82 (br s, 1H), 8.00 – 7.98 (m, 2H), 7.74 (d, 1H, *J* = 4.1 Hz), 7.69 (dd, 1H, *J* = 8.2 and 1.6 Hz), 7.14 (d, 1H, *J* = 8.1 Hz), 1.99 – 1.83 (m, 7H) and 1.40 – 1.37 (m, 3H); MS ((-)-APCI) *m/z* 323 [M-H]⁻.

Example 36**5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indole]-2''(1''H)-thione****5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one:** Prepared

5 according to the procedure for Example 5: mp 171-172 °C; ¹H-NMR (CDCl₃) δ 8.43 (s, 1H), 7.62 (d, 1H, *J* = 1.8 Hz), 7.42 (dt, 1H, *J* = 6.2, 2.0 Hz), 7.39 - 7.37 (m, 1 H), 7.33 (dt, 1H, *J* = 5.1, 1.3 Hz), 7.26 (dq, 1 H, *J* = 5.9, 2.1 Hz), 7.05 - 6.99 (m, 2H), 2.03 - 1.64 (m, 10H); MS ((+)APCI) *m/z* 296 [M+H]⁺.

The title compound was prepared from 5'-(3-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (0.70 g) and phosphorous pentasulfide (0.4 g) in toluene (20 ml) at reflux. After 3 h. the reaction was cooled, poured into sat. aqueous sodium hydrogen carbonate solution, and extracted with EtOAc, the organic layer was separated, dried (MgSO₄) and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc-hexane gradient elution) to afford the product (0.42 g) as an off white solid: ¹H NMR (DMSO-d₆) δ 12.75 (br s, 1H), 7.95 (d, 1H, *J* = 1.5 Hz), 7.64 (dd, 1H, *J* = 8.13 and 1.5 Hz), 7.53 - 7.48 (m, 3H), 7.21 - 7.14 (m, 2H), 1.99 - 1.83 (m, 7H) and 1.40 - 1.37 (m, 3H); MS ((-)-APCI) *m/z* 310 [M-H]⁻.

Example 37**5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione****5'-(3-Hydroxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one:** Prepared

20 according to the procedure for example 5: mp. 213 - 216 °C; ¹H NMR (CDCl₃) δ 1.60-1.96 (m, 10H), 6.78 - 6.82 (m, 1H), 6.94 (d, 1H, *J* = 8 Hz), 7.01 - 7.04 (m, 2H), 7.23 (t, 1H, *J* = 7.7 Hz), 7.38 (d, 1H, *J* = 8 Hz), 7.61 (s, 1H), 8.91 (s, 1H) and 9.73 (s, 1H, 25 br); MS ((+)-APCI) *m/z* 294 [M+H]⁺.

The title compound was prepared from 5'-(3-hydroxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one (100 mg) and Lawesson's reagent (110 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.0045 g) as an off white solid: ¹H NMR (CDCl₃) δ 9.59 (br s, 1H), 7.89 (s, 1H), 7.49 (dd, 1H, *J* = 8.1 and 1.5 Hz), 7.33 (t, 1H, 30

$J = 7.9$ Hz), 7.15 – 7.10 (m, 3H), 6.84 (dd, 1H, $J = 8.0$ and 2.2 Hz), 2.17 – 2.05 (m, 2H), 1.98 – 1.88 (m, 5H) and 1.57 – 1.53 (m, 3H): MS ((-)-APCI) m/z 308 [M-H].

Example 38

5 5-(3-chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione

A solution of oxindole (40 g, 0.3 mol) in dry THF (400 ml) under N_2 was cooled to $-25^\circ C$ and treated drop wise with *n*-butyl lithium (2.5M in hexanes, 240 ml, 0.6 mol). To the resulting solution was added N,N,N',N' -tetramethylethylenediamine (90.4 ml, 0.6 mol). After 30 min. iodoethane (48 ml, 0.6 mol) was added and the
10 reaction mixture was allowed to warm to room temperature and stirred over night. The reaction mixture was poured into aqueous NH_4Cl solution, extracted with EtOAc (2x) and the combined organic layers were washed with dil. HCl, water, brine, dried ($MgSO_4$) and concentrated. The residual oil was triturated with hexane to afford the crude product (24.5 g, 51%). A sample (3 g) was recrystallized from EtOAc/hexane
15 to obtain 3-ethyl-indol-2-one (1.4 g), m.p. $100 - 101^\circ C$; 1H -NMR ($DMSO-d_6$) δ 0.76 (t, 3H, $J = 7.5$ Hz), 1.8 – 2.0 (m, 2H), 3.38 (t, 3H, $J = 5.7$ Hz), 6.8 (dt, 1H, $J = 7.69$, 0.45 Hz), 6.93 (dt, 1H, $J = 7.45$, 1.10 Hz), 7.15 (m, 1H), 7.22 (m, 1H), 10.3 (s, 1H); MS (ESI) m/z 270 [M+H].

A solution of 3-ethyl-indol-2-one (16 g, 0.1 mol) in dry THF (200 ml) under
20 N_2 was cooled to $-25^\circ C$ and treated drop wise with *n*-butyllithium (2.5M in hexanes, 80 ml, 0.2 mol). To the resulting solution was added N,N,N',N' -tetramethylethylenediamine (30 ml, 0.2 mol). After 30 min. iodoethane (8 ml, 0.1 mol) was added and the reaction mixture was allowed to warm to RT and stirred over night. The reaction mixture was poured into an aqueous NH_4Cl solution, extracted
25 with EtOAc (2x) and the combined organic layers were washed with dil. HCl, water, brine, dried ($MgSO_4$) and concentrated. The residual oil was triturated with hexane to afford 3,3-diethylindol-2-one (9 g, 45%), m.p. $156 - 159^\circ C$; 1H NMR ($DMSO-d_6$) δ 10.44 (s, 1H), 7.70 - 7.69 (t, 1H), 7.62 - 7.59 (m, 1H), 7.58 (d, 1H $J = 1.7$ Hz), 7.53 - 7.50 (m, 1H), 7.45 - 7.41 (t, 1H), 7.36 - 7.35 (m, 1H), 7.34 - 7.33 (m, 1H), 6.91 - 6.89 (d, 1H

- 88 -

$J=8.2\text{Hz}$), 1.87 - 1.80 (m, 2H), 1.77 - 1.70 (m, 2H), 0.54 - 0.50 (t, 6H); MS (+ESI) m/z 190 (M+H).

A solution of 3,3-diethylindol-2-one (8 g, 40 mmol) and sodium acetate (4 g, 48 mmol) in acetic acid (100 ml) was treated with bromine (6.4 g, 40 mmol). After
5 30 min. the mixture was diluted with water and extracted with EtOAc (2x); the combined organic layers were washed with water, sat. sodium hydrogen carbonate solution, then brine, dried (MgSO_4) and evaporated to afford the crude product (7.6 g, 75%). A sample was recrystallized from EtOAc/hexane to obtain 5-bromo-1,3-dihydro-3,3-diethyl-[2H]-indol-2-one, m. p. 164 - 165 °C; $^1\text{H-NMR}$ (DMSO-d_6) δ
10 10.45 (s, 1H), 7.41-7.40(d, 1H, $J = 2.2\text{Hz}$), 7.34 - 7.31 (m, 1H), 6.78 - 6.76 (d, 1H $J = 8.2\text{ Hz}$), 1.78-1.65 (m, 4H), 0.50 - 0.46 (m, 6H); MS (-ESI) m/z 266/268 (M-H).

A solution of 5-bromo-1,3-dihydro-3,3-diethyl-[2H]-indol-2-one (2.7 g, 10 mmol), 3-chlorophenylboronic acid (1.6 g, 10 mmol), potassium carbonate (4 g, 30 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.5 g, 0.4 mmol) in
15 dimethoxyethane (100 ml), ethanol (25 ml), and water (25 ml) was heated to reflux for 6 hours. After cooling to room temperature, the mixture was diluted with water and extracted with EtOAc (2x). The combined organic extracts were washed with water, then brine, dried (MgSO_4) and evaporated. The residue was purified by column chromatography (SiO_2 , EtOAc:hexane 1:3) to afford 5-(3-chloro-phenyl)-3,3-diethyl-1,3-dihydro-indol-2-one compound (0.8 g, 27%), m.p. 195 - 197 °C; $^1\text{H-NMR}$ (DMSO-d_6) δ 7.70 (t, 1H, $J = 2\text{ Hz}$), 7.62 - 7.60 (m, 1H), 7.58 (d, 1H, $J = 1.7\text{ Hz}$),
20 7.52, (dd, 1H, $J = 8.1, 2\text{ Hz}$), 7.43 (t, 1H, 7.9 Hz), 7.36 - 7.33 (m, 1H), 6.90 (d, 1H, $J = 8.1\text{ Hz}$), 1.87 - 1.70 (m, 4H) and 0.52 (t, 6H, $J = 7.4\text{ Hz}$); MS (+APCI) m/z 300/302 (M-H).

25 The title compound was prepared from 5-(3-chloro-phenyl)-3,3-diethyl-1,3-dihydro-indol-2-one compound (100 mg) and Lawesson's reagent (100 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.023 g) as a yellow solid: $^1\text{H NMR}$ (DMSO-d_6) δ 12.73 (br s, 1H), 7.77 (t, 1H, $J = 1.8\text{ Hz}$), 7.75 (d, 1H, $J = 1.6\text{ Hz}$), 7.68 - 7.62 (m, 2H), 7.48 (t, 1H, $J = 7.9\text{ Hz}$), 7.40 (d, 1H, $J = 8.3$

- 89 -

Hz), 7.09 (d, 1H, $J = 8.1$ Hz), 2.07 – 2.00 (m, 2H), 1.86 – 1.79 (m, 2H) and 0.37 (t, 6H, $J = 7.3$ Hz): MS ((-)-APCI) m/z 314/316 [M-H]⁻.

Example 39

5 5-[4-Fluoro-3-(trifluoromethyl)phenyl]spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione

5-[4-Fluoro-3-(trifluoromethyl)phenyl]spiro[cyclohexane-1,3-[3H]indol]-2(1H)-one was prepared from (2'-oxo-2,3-dihydrospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)boronic acid (2.5 g, 10 mmol) and 5-bromo-2-fluoro-trifluoromethylbenzene (2 g, 8 mmol) as described for Example 5, to afford the title compound (0.87 g, 30%) as a solid: mp. 222 °C; ¹H NMR (DMSO-d₆) δ 1.5 - 1.8 (m, 8 H), 1.8 - 2.0 (m, 2 H), 6.92 (d, 1 H, $J = 8.13$ Hz), 7.51 (dd, 1 H, $J = 8.13, 1.76$ Hz), 7.55 (dd, 1 H, $J = 10.54, 9.01$ Hz), 7.72 (d, 1 H, $J = 1.76$ Hz), 7.90 (dd, 1 H, $J = 7.03, 2.20$ Hz), 7.98 (m, 1 H) and 10.39 (s, 1 H); MS (EI) m/z 363 (M⁺).

15 The title compound was prepared from 5-[4-Fluoro-3-(trifluoromethyl)phenyl]spiro[cyclohexane-1,3-[3H]indol]-2(1H)-one (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.016 g) as a yellow solid: ¹H NMR (DMSO-d₆) δ 12.75 (br s, 1H), 8.06 – 8.00 (m, 1H), 7.96 – 7.92 (m, 2H), 7.66 – 7.56 (m, 2H), 7.16 (d, 1H, $J = 8.1$ Hz), 1.99 – 1.83 (m, 7H) and 1.41 – 1.38 (m, 3H): MS ((-)-APCI) m/z 378 [M-H]⁻.

Example 40

4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile

25 The title compound was prepared from 4-(1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the title compound (0.050 g) as an orange solid: ¹H NMR (DMSO-d₆) δ 12.80 (br s, 1H), 8.04 (d, 1H, $J = 1.3$ Hz), 7.98 (t, 1H, $J = 7.5$ Hz), 7.92

- 90 -

(dd, 1H, $J = 11.3$ and 1.3 Hz), 7.76 (d, 2H, $J = 8.0$ Hz), 7.18 (d, 1H, $J = 8.2$ Hz), 1.99 – 1.82 (m, 7H) and 1.40 – 1.38 (m, 3H); MS ((-)-APCI) m/z 335 [M-H].

Example 41

5 **5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H] indol]-5-yl) 4-n-butyl-2-thiophenecarbonitrile**

The title compound was prepared in a manner similar to Example 5 from 5-bromo-4-n-butyl thiophenecarbonitrile (1.24 g, 5.1 mmol), (1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H] indol]-5-boronic acid (1.24 g, 5.05 mmol),
10 tetrakis(triphenylphosphine) palladium (0.25 g), potassium carbonate (2.75 g, 21 mmol), water (10 mL), and dimethoxyethane (50 mL) heated at reflux for 5 hours to afford 5-(1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H] indol]-5-yl) 4-n-butyl-2-thiophenecarbonitrile (1 g, 54%), m.p. 130-132° C. ^1H NMR (DMSO- d_6) δ 10.56 (s, 1H), 7.92 (s, 1H), 7.52-7.51 (d, 1H, $J = 1.2$ Hz), 7.32-7.29 (dd, 1H, $J = 1.5$ Hz), 6.98
15 - 6.96 (d, 1H, $J = 8.0$ Hz), 2.64 - 2.59 (t, 2H), 1.99 - 1.86 (m, 2H), 1.70 - 1.50 (m, 11 H), 1.32 - 1.22 (m, 2H), 0.86 - 0.82 (t, 3H); MS (APCI (+)) m/z 365 [M+H] $^+$; IR (KBr) 1620, 1700; 2200 cm^{-1} ; Anal. $\text{C}_{22}\text{H}_{24}\text{N}_2\text{OS} \cdot 1/4 \text{H}_2\text{O}$. calc. C, 71.61; H, 6.69; N 7.59. observed C, 71.13; H, 6.61; N, 6.91.

The title compound was prepared from 5-(1,2-dihydro-2-oxospiro[cyclohexane-1,3-[3H] indol]-5-yl) 4-n-butyl-2-thiophenecarbonitrile (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.050 g) as an orange solid: ^1H NMR (DMSO- d_6) δ 12.83 (br s, 1H), 7.95 (s, 1H), 7.77 (s, 1H), 7.44 (d, 1H, $J = 8.1$ Hz), 7.18 (d, 1H, $J = 8.1$ Hz), 2.63 (t, 1H, $J = 8.0$ Hz), 1.99 – 1.77 (m, 7H), 1.60 – 1.50
20 (m, 2H), 1.39 – 1.35 (m, 3 H), 1.29 – 1.22 (m, 2H) and 0.81 (t, 3H, 7.3 Hz); MS ((-)-APCI) m/z 379 [M-H].

Example 42

30 **5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indole]-2(1H)-thione**
The title compound was prepared from 5-(3-Fluoro-5-methoxyphenyl)spiro

- 91 -

[cyclohexane-1,3-[3H]indole]-2(1H)-one (90 mg) and Lawesson's reagent (90 mg) in toluene (10 ml) at reflux, according to General Procedure A, to afford the product (0.043 g) as an off-white solid: ¹H NMR (DMSO-d₆) δ 12.74 (br s, 1H), 7.90 (s, 1H), 7.63 (dd, 1H, *J* = 8.1 and 1.2 Hz), 7.13 (d, 1H, *J* = 8.1 Hz), 7.08 (d, 1H, *J* = 10 Hz), 7.01 (s, 1H), 6.83 (dt, 1H, *J* = 11 and 2.0 Hz), 1.99 – 1.83 (m, 7H) and 1.40 – 1.37 (m, 3H); MS ((-)-APCI) *m/z* 340 [M-H].

Example 43

5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine

10 To a solution of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indole]-2'(1H)-thione (0.74 g, 2.25 mmol) in dry THF (15 ml) was added sodium hydride (60% in oil, 0.1 g, 2.5 mmol) at room temperature. After 15 min., methyl iodide (0.18 ml, 2.88 mmol) was added. After 1h, the reaction mixture was partitioned between water and EtOAc, the organic layer was washed with brine, dried (MgSO₄) and
15 evaporated to give 5-(3-chlorophenyl)-2-(methylthio)spiro[cyclohexane-1,3'-[3H]indole] (0.80 g, 100%) which was used without further purification:

To a solution of the last cited compound (1.96 g, 5.73 mmol) in DMSO (20 ml) was added hydroxylamine (60% in water, 5 ml) and the mixture was heated to 120 °C. After 1h., the reaction was cooled, partitioned between diethyl ether and
20 saturated aqueous ammonium chloride solution. The organic layer was washed with water and brine and then dried (MgSO₄) and evaporated. The crude product was then crystallized from MeOH to afford the title compound (1.67 g, 5.08 mmol, 89%) as a white solid: ¹H NMR (CDCl₃) δ 7.52 (t, 1H, *J* = 1.7 Hz), 7.43 – 7.28 (m, 7H), 6.83 (d, 1H, *J* = 8 Hz) and 1.98 – 1.51 (m, 10H); MS (ESI (+)) *m/z* 327/329 [M+H]⁺.

25

Example 44

N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine

To a solution of 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine (0.23 g, 0.71 mmol) in methylene chloride-methanol (9:1, 10 ml)
30 was added acetic anhydride (0.08 ml, 0.8 mmol) and 4-dimethylaminopyridine

(catalytic amount) under a nitrogen atmosphere. After 20 min., the reaction was evaporated and the product purified by column chromatography (SiO₂, methanol: methylene chloride 5:95). The product was then triturated with di-iso-propylether to afford the title compound (0.12 g, 0.32 mmol, 45%): ¹H NMR (CDCl₃) δ 7.52 – 7.51 (m, 2H), 7.43 – 7.27 (m, 5H), 6.88 (d, 1H, J = 8 Hz), 2.27 (s, 3H), 2.04 – 1.92 (m, 4H), 1.84 – 1.74 (m, 4H) and 1.72 – 1.57 (m, 2H); MS (ESI (+)) m/z 369/371 [M+H]⁺; C₂₁H₂₁ClN₂O₂. 0.5 H₂O requires C 66.98: H 5.64: N 7.34. Found C 66.74: H 5.86: N 7.41.

10

Example 45**5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol-2'-(1'H)-one oxime**

Prepared according to the method for Example 43 from 5'-(3-fluorophenyl) spiro[cyclohexane-1,3'-[3H]indole]-2'-(1'H)-thione (0.59 g, 1.90 mmol) to afford the title compound (0.053 g, 0.17 mmol, 10%): ¹H NMR (DMSO-d₆) δ 9.59 (s, 1H), 9.40 (s, 1H), 7.57 (d, 1H, J = 1.5 Hz), 7.46 – 7.39 (m, 4H), 7.11 – 7.05 (m, 1H), 6.80 (d, 1H, J = 8.1 Hz), 2.04 – 1.97 (m, 2H), 1.82 – 1.74 (m, 2H) and 1.66 – 1.42 (m, 6H): MS (ESI (-)) m/z 309 [M-H]⁻; C₁₉H₁₉FN₂O requires C: 73.53, H: 6.17, N: 9.03. Found C: 73.33, H: 6.07, N: 8.83.

20

Example 46**5'-(2-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol-2'-(1'H)-one oxime**

5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O- benzyloxime). 5'-Bromo-2'-(methylthio)spiro[cyclohexane-1,3'-[3H]indole] (9.0 g, 28.98 mmol) and O-benzylhydroxylamine hydrochloride (13.8 g, 86.9 mmol) were combined in methanol (150 mL) and heated to 45°C for 6 hours. Methanol was evaporated in vacuo. Ethyl acetate was added to the residue and this mixture was washed with ammonium chloride solution. Ethyl acetate was dried over magnesium sulfate, ethyl acetate collected evaporated in vacuo and the residue was flash chromatographed on alumina 90 (9:1 Hexane/EtOAc) to the desired product (6.5 g, 60%).

- 93 -

¹H NMR (DMSO-d₆, 300 MHz) δ 1.38-1.70 (m, 8H), 1.92-2.06 (m, 2H), 5.06 (s, 2H), 6.71 (d, 1H, J = 8.26 Hz), 7.22-7.43 (m, 7H), 9.62 (s, 1H).

Procedure A

5 5'-(2-Fluorophenyl)-spiro[cyclohexane-1,3'-[3H]indol]-2'(1H)-one 2(O-benzyloxime). 5'-Bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (1.0 g, 2.6 mmol), and tetrakis(triphenyl phosphine) Pd (0) (0.14 g, 0.12 mmol) were stirred under an atmosphere of nitrogen in ethylene glycol dimethyl ether (23 mL). After 15 minutes, 2-fluorophenyl boronic acid (0.72 mg, 5.2 mmol) was
10 added, followed by sodium carbonate (1.6 g, 15.6 mmol) in water (6.0 mL). The reaction was heated to reflux overnight, cooled to room temperature and filtered through a Celite plug. Saturated ammonium chloride was added. The water layer was extracted with ethyl acetate (3 X 100 mL). The combined organic layers were dried (MgSO₄), filtered, and the solvent removed in vacuo. The product was purified by
15 flash silica gel chromatography; (eluant: 10:0.5 hexane: ethyl acetate) to give the **desired target compound** (0.75 g, 1.8 mmol, 72%) as a viscous oil. ¹H NMR (500 MHz, DMSO-d₆) δ 1.44 - 1.73 (8H, m) 1.93 - 2.06 (2H, q) 5.00 (2H, s) 6.88 (1H, d, J = 8.1 Hz) 7.24 - 7.38 (6H, m) 7.44 - 7.56 (5H, m) 9.64 (1H, s); MS (ESI(+ ve)) m/z 399 (M-H)⁺.

Procedure B

20 A solution of 5'-(2-Fluorophenyl)-spiro[cyclohexane-1,3'-[3H]indol]-2'(1H)-one 2(O-benzyloxime) (0.55 g, 1.37 mmol) in ethanol (15 mL) was added to Palladium on carbon (10%, 0.11 g) in ethanol (10 mL). The mixture was stirred under an atmosphere of hydrogen (balloon) for 24 h at room temperature. The reaction
25 mixture was filtered through a Celite plug and the filtrate was concentrated in vacuo. The product was purified by flash silica gel chromatography (hexane: ethyl acetate, gradient elutions) to give the **title compound** (0.45 g, 1.12 mmol, 82%), mp. 200 – 203 °C; ¹H NMR (500 MHz, DMSO-d₆) δ 1.45 - 1.73 (8H, m) 1.96 - 2.00 (2H, q) 6.83 (1H, d, J = 7.9) 7.23 - 7.50 (6H, m) 9.42 (1H, s) 9.58 (1H, s); MS (ESI(+ ve)) m/z 311 (M+H)⁺.

Example 47**5'-(4-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol-2'(1'H)-one oxime**

5'-(4-Fluorophenyl)-spiro[cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime). Prepared from 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (1.0 g, 2.6 mmol), and 4-fluorophenyl boronic acid (0.72 g, 5.2 mmol) according to Example 46 procedure A. The product was purified by flash silica gel chromatography; (eluant: 10:0.5 hexane: ethyl acetate) to give the desired product (0.70 g, 1.7 mmol, 67 %) as a viscous oil. ¹H NMR (500 MHz, DMSO-d₆) δ 1.42 - 1.77 (8H, m) 1.95 - 1.99 (2H, q) 5.00 (2H, s) 6.84 (1H, d, J= 8.1 Hz) 7.21 - 7.63 (1H, m) 9.58 (1H, s); MS (ESI(- ve)) m/z 399 (M-H).

The product was synthesized using 5'-(4-Fluorophenyl)-spiro[cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (0.70 g, 1.74 mmol), according to Example 45 procedure B. The product was purified by flash silica gel chromatography; (hexane: ethyl acetate, gradient elution) to give the title compound (0.44 g, 1.4 mmol, 81 %), Mp. 205 - 208°C; ¹H NMR (500 MHz, DMSO-d₆) δ 1.43 - 1.77 (8H, m) 2.00 - 2.05 (2H, q) 6.80 (1H, d, J= 8.2 Hz) 7.21 - 7.24 (2H, m) 7.33 - 7.35 (1H, dd, J= 1.9 Hz) 7.49 (1H, s) 7.60 - 7.63 (2H, m) 9.35 (1H, s) 9.56 (1H, s); MS (ESI(+ ve)) m/z 311 (M+H)⁺.

20

Example 48**5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol-2'(1'H)-one oxime**

5'-(3,4-Difluorophenyl)-spiro[cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime). Prepared from 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (1.0 g, 2.6 mmol) and 3,4 -difluorophenyl boronic acid (1.6 g, 5.2 mmol of a 50 % solution of acid in THF/water) according to Example 46 procedure A. The product was purified by flash silica gel chromatography (eluant: 10:0.5 hexane: ethyl acetate) to give the desired product (0.75 g, 1.7 mmol, 69 %) as a viscous oil. ¹H NMR (500 MHz, DMSO-d₆) δ 1.41 - 1.78 (8H, m) 1.95 - 1.99 (2H,

- 95 -

q) 5.00 (2H, s) 6.82 (1H, d) 7.28 – 7.46 (8H, m) 7.58 (1H, q) 7.67 – 7.71 (1H, m) 9.61 (1H, s); MS (ESI(- ve)) m/z 417 (M-H).

Reaction of the last cited compound (0.70 g, 1.6 mmol) according to Example 45 procedure B, afforded the title compound (0.44 g, 1.3 mmol, 80 %), ¹H NMR (500 MHz, DMSO- d_6) δ 1.42 – 1.79 (8H, m) 2.01 – 2.05 (2H, q) 6.78 – 6.80 (1H, d) 7.39 – 7.46 (3H, m) 7.55 (1H, s) 7.70 (1H, m) 9.10 (1H, s) 9.59 (1H, s); MS (ESI(+ ve)) m/z 329 (M+H)⁺.

Example 49

10 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol-2'(1'H)-one oxime

5'-(3-Methoxyphenyl)-spiro[cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime). Prepared from 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (1.0 g, 2.6 mmol) and 3-methoxyphenyl boronic acid (0.79 g, 5.2 mmol) according to Example 46 procedure A. The product was purified by flash silica gel chromatography; (eluant: 10:0.5 hexane: ethyl acetate) to give the desired product (0.80 g, 1.9 mmol, 75 %) as a viscous oil. ¹H NMR (500 MHz, DMSO- d_6) δ 1.43 - 1.78 (8H, m) 1.95 - 2.00 (2H, q) 3.80 (3H, s) 5.00 (2H, s) 6.82 - 6.86 (2H, m) 7.10 - 7.16 (2H, m) 7.28 - 7.53 (10H, m) 9.57 (1H, s); MS (ESI(- ve)) m/z 411 (M-H).

Reaction of the last cited compound (0.80 g, 1.9 mmol) according to Example 20 46 procedure B, afforded the title compound (0.48 g, 1.4 mmol, 77 %), as a white solid. Mp. 101 - 104°C; ¹H NMR (500 MHz, DMSO- d_6) δ 1.44 - 1.78 (8H, m) 1.99 - 2.03 (2H, q) 3.81 (3H, s) 6.78 (1H, d) 6.85 (1H, d) 7.10 - 7.16 (2H, m) 7.30 - 7.38 (2H, m) 7.50 (1H, d) 9.35 (1H, s) 9.56 (1H, s); MS (ESI(+ ve)) m/z 323 (M+H)⁺.

25 Example 50

5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol-2'(1'H)-one oxime

5'-(3-Nitrophenyl)-spiro[cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime). Prepared from 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (1.0 g, 2.6 mmol) and 3-Nitrophenyl boronic acid (0.86 g, 5.2 mmol) according to Example 46 procedure A. Purification by flash silica gel 30

- 96 -

chromatography (eluant : 10:0.5 hexane: ethyl acetate) afforded the desired compound (0.60 g, 1.4 mmol, 55 %) as a viscous oil. ¹H NMR (500 MHz, DMSO-d₆) δ 1.42 - 1.82 (8H, m) 2.02 - 2.04 (2H, q) 5.01 (2H, s) 6.88 (1H, d) 7.28-7.71 (8H, m) 8.08 - 8.13 (2H, m) 8.38 (1H, d) 9.69 (1H, s); MS (ESI(- ve)) *m/z* 426 (M-H)⁻.

5

Procedure C

The last cited compound (0.54 g, 1.26 mmol) was dissolved in dry methylene chloride (25 mL) and cooled to -78 °C under nitrogen. Boron tribromide (3.8 mL, 3.8 mmol, 1.0 M in methylene chloride) was added drop-wise over 5 minutes. After 30 minutes the reaction was quenched with saturated sodium bicarbonate (5 mL). The reaction mixture was allowed to warm to room temperature, the layers were separated and the aqueous layer was extracted with methylene chloride. The combined organic layers were dried (Na₂SO₄), filtered, and the solvent removed *in vacuo*. The product was purified by flash silica gel chromatography (eluant: 8:1 hexane: ethyl acetate) to give afford the **title compound** (0.33 g, 0.9 mmol, 78 %). Mp. 221 - 224 °C; ¹H NMR (500 MHz, DMSO-d₆) δ 1.42 - 1.83 (8H, m) 1.99 - 2.07 (2H, q) 6.84 - 6.85 (1H, dd) 7.50 - 7.52 (1H, m) 7.67 - 7.71 (2H, m) 8.08 - 8.12 (2H, m) 8.37 - 8.38 (1H, d) 9.48 (1H, s) 9.64 (1H, s); MS (ESI(+ ve)) *m/z* 338 (M+H)⁺.

15

Example 51

20

5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol-2'(1'H)-one oxime

25

3-[Spiro[cyclohexane-1,3'-[3H]indol]-(1'H)-one-2'-(O-

benzyloxime)]benzonitrile [3H]indol]-5-yl]benzonitrile. Prepared from 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O- benzyloxime) (1.0 g, 2.6 mmol) and 3-cyanophenyl boronic acid (0.76 g, 5.2 mmol) according to Example 46 procedure A. The product was purified by flash silica gel chromatography (eluant: 10:0.5 hexane: ethyl acetate) to give the desired **product** (0.75 g, 1.8 mmol, 71 %) as a viscous oil. ¹H NMR (500 MHz, DMSO-d₆) δ 1.41 - 1.81 (8H, m) 1.96 - 2.03 (2H, q) 5.01 (2H, s) 6.86 (1H, d) 7.28 - 7.33 (9H, m) 7.95 - 7.97 (1H, d) 8.12 (1H, s) 9.65 (1H, s); MS (ESI(- ve)) *m/z* 406 (M-H)⁻.

- 97 -

Reaction of the last cited compound (0.17 g, 0.43 mmol) and boron tribromide (1.2 mL, 1.2 mmol) according to Example 50 procedure C afforded the **title compound** (0.06 g, 0.2 mmol, 47 %) as a white solid, Mp. 198 – 200°C; ¹H NMR (500 MHz, DMSO-d₆) δ 1.41 – 1.80 (8H, m) 1.97 – 2.04 (2H, q) 6.80 (1H, q) 7.45 – 7.69 (4H, m) 7.93 – 7.95 (1H, dd) 8.10 (1H, s) 9.42 (1H, s) 9.59 (1H, s); (ESI(+ ve)) *m/z* 318 (M+H)⁺.

Example 52

3-[1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3']-[3H]indol]-5'-yl]-5-fluorobenzonitrile

To a solution of 3-fluoro-5-cyano-bromobenzene (0.4 g, 2.0 mmol) in dry DMF (10 ml) was added diboron pinacolate ester (0.63 g, 2.5 mmol), potassium acetate (0.65 g, 6.7 mmol) and PdCl₂ (dppf) (0.2. g) and the reaction was heated to 80 °C under a nitrogen atmosphere. After 8 h. from 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O- benzyloxime) (0.2 g, 0.5 mmol), PdCl₂ (dppf) (0.05 g) and sodium carbonate (1.30 g, 12.5 mmol) were added and heating at 80 °C was continued. After 8h. the reaction was cooled and partitioned between water and ethyl acetate, the organic layer was washed with brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography (SiO₂, EtOAc:hexane 1:20) to give the desired product (0.14 g, 0.33 mmol, 66%).

Reaction of the last cited compound (0.14 g, 0.33 mmol) and boron tribromide (1.0 ml, 1.0 mmol) according to Example 50 procedure C afforded the title compound (0.019 g, 0.05 mmol, 17%): ¹H NMR (300 MHz, DMSO-d₆) δ 9.65 (s, 1H), 9.49 (s, 1H), 8.04 (m, 1H), 7.89 (dt, 1H, *J* = 10.5 and 2 Hz), 7.72 – 7.68 (m, 2H), 7.54 (d, 1H, *J* = 8.1 Hz), 6.80 (d, 1H, *J* = 8.1 Hz), 2.05 – 1.99 (m, 2H), 1.84 – 1.76 (m, 2H) and 1.65 – 1.44 (m, 6H): MS (ESI(+ ve)) *m/z* 336 (M+H)⁺.

Example 53**5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile**

5 **4-Methyl-5-trimethylstannanyl-thiophene-2-carbonitrile**. Prepared from 5-bromo-4-methyl-thiophene-2-carbonitrile (3.08 g, 15.2 mmol), tetrakis(triphenyl phosphine) Pd (0) (0.82 g, 0.71 mmol), hexamethylditin (5.0 g, 15.2 mmol) and ethylene glycol dimethyl ether (20 mL) under nitrogen. The mixture was heated to reflux for 14 hours. The reaction mixture was concentrated in vacuo and purified
10 using flash silica gel chromatography (eluant: 2% MeOH: methylene chloride) to recover the desired **product** (2.8 g, 0.01 mmol, 67 %) as a runny oil. ¹H NMR (300 MHz, DMSO-d₆) δ 0.41 (9H, s), 2.28 (3H, s), 7.83 (1H, s).

The last cited compound (0.20 g, 0.50 mmol), dichlorobis
(triphenylphosphine) palladium(II) (0.02 g, 0.03 mmol) and triphenylarsine (0.03 g,
15 0.13 mmol) in DME (8.0 mL) were stirred under nitrogen for 20 minutes. 5'-Bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O- benzyloxime) (0.18 g, 0.64 mmol) was added in a solution of DME (2.0 mL). The solution was heated to reflux overnight. The reaction solution was concentrated in vacuo and purified by flash silica gel chromatography (eluant 12:1 hexane : ethyl acetate) to give the crude
20 product(0.10 g, 0.25 mmol, 50%) which was used without further purification

Boron tribromide (2.6 mL, 2.6 mmol of a 1.0 M solution in methylene chloride) was added to a solution of the last product (0.37 g, 0.86 mmol) in dry methylene chloride (1.7 mL) at -78°C. The solution was stirred for 30 minutes and quenched with saturated sodium bicarbonate (10 mL). The mixture was allowed to
25 warm to room temperature and the layers were separated. The organic layer was dried (Na₂SO₄), filtered and concentrated in vacuo to give crude product which was purified by flash silica gel chromatography (eluant: 6:1 hexane: ethyl acetate) to give the title compound (0.02 g, 24 %): Mp. 173-176 °C; ¹H NMR (500 MHz, DMSO-d₆) δ 1.44 - 1.73 (8H, m), 1.96 - 2.00 (2H, m), 2.28 (3H, s), 6.82 - 6.84 (1H, m), 7.24 - 7.26 (1H,

- 99 -

dd, $J = 1.7$ Hz), 7.38 (1H, m) 7.82 (1H, m) 9.51 (1H, m) 9.66 (1H, m); MS (ESI(+ ve)) m/z 338 (M+H)⁺.

Example 54

5 **5-(Spiro[cyclohexane-1,3'-[3H]indole]-2'(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile**

10 To a solution of 2-cyanothiophene (1.0 g, 9.16 mmol) and tri-*iso*-propylborate (2.3 ml, 10 mmol) in dry THF (30 ml) under nitrogen at -78 °C was added, dropwise, lithium hexamethyldisilazide (1M in THF, 10 ml, 10 mmol). After 30 min., the reaction was quenched with 1N HCl, then extracted with ethyl acetate, the organic layer was washed with water, dried (Na₂SO₄) and evaporated to the product (1.25g, 8.17 mmol, 89%) which was used without further purification: ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.75 (br s, 2H), 7.97 (d, 1H, $J = 8$ Hz) and 7.73 (d, 1H, $J = 8$ Hz): MS (ESI(- ve)) m/z 152 (M-H)⁻.

15 Prepared from the last cited product (0.91 g, 5.95 mmol) and 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O- benzyloxime) (1.53 g, 3.97 mmol) according to Example 46 procedure A. Purification by flash silica gel chromatography (eluant: 5: 1 hexane: THF) gave the desired **product** (0.66 g, 1.59 mmol) which was used without further purification: MS (ESI(- ve)) m/z 412 (M-H)⁻.

20 Reaction of the last cited compound (0.60 g, 1.45 mmol) and boron tribromide (1M in dichloromethane, 5 mL, 5 mmol) according to Example 50 procedure C afforded the **title compound** (0.036 g, 0.11 mmol, 8 %): ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.71 (s, 1H), 9.62 (s, 1H), 7.92 (d, 1H, $J = 3.9$ Hz), 7.63 (d, 1H, $J = 1.5$ Hz), 7.54 (d, 1H, $J = 3.9$ Hz), 7.47 (dd, 1H, $J = 8.1$ and 1.6 Hz), 6.78 (d, 1H, $J = 8.1$ Hz), 2.13 – 1.90 (m, 2H) and 1.78 – 1.60 (m, 6H): MS (ESI(+ ve)) m/z 324 (M+H)⁺.

25

Example 55

4-(Spiro[cyclohexane-1,3'-[3H]indole]-2'(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile

4-(Trimethylstannyl)-2-thiophenecarbonitrile A solution of 3-bromo-2-

5 thiophenecarbonitrile (0.8 g, 4.3 mmol), tetrakis(triphenylphosphine)palladium(0) (0.25 g, 0.2 mmol) and hexamethylditin (1.4 g, 4.3 mmol) in dimethoxyethane (5 cm³) was heated under reflux for 14 h then cooled to RT. The reaction mixture was absorbed onto florisil and purified by column chromatography (SiO₂, methylene chloride: hexane 1:9) to afford the subtitled compound (1.04 g, 3.8 mmol, 90%) as a
10 clear viscous oil: ¹H NMR (CDCl₃) δ 0.35 (s, 9H), 7.56 (d, *J* = 0.9 Hz, 1H), 7.66 (d, *J* = 0.9 Hz, 1H).

To a solution of 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O- benzyloxime) (1.65 g, 4.28 mmol), 4-(trimethylstannyl)-2-thiophenecarbonitrile (1.48 g, 5.44 mmol), triphenylarsine (330 mg) in dry dimethoxy ethane (20 ml), under
15 a nitrogen atmosphere was added bis(triphenylphosphine)palladium (II) chloride, and the mixture was heated under reflux for 16 h. After cooling to room temperature the mixture was evaporated, and the residue purified by column chromatography (SiO₂, EtOAc:hexane, gradient elution) to afford the desired product (0.61 g, 1.47 mmol, 56%).

20 Reaction of the last cited compound (0.61 g, 1.47 mmol) and boron tribromide (1M in dichloromethane, 4.5 mL, 4.5 mmol) according to Example 50 procedure C afforded the **title compound** (0.084 g, 0.26 mmol, 18 %): ¹H NMR (300 MHz, DMSO-d₆) δ 9.61 (s, 1H), 9.42 (s, 1H), 8.41 (s, 1H), 8.18 (s, 1H), 7.65 (s, 1H), 7.48 (dd, 1H, *J* = 8.1 and 0.9 Hz), 6.76 (d, 1H, *J* = 8.1 Hz), 2.03 – 1.96 (m, 2H) and
25 1.78 – 1.42 (m, 6H): MS (ESI(+ ve)) *m/z* 324 (M+H)⁺.

Example 56

5-(Spiro[cyclohexane-1,3'-[3H]indole]-2'(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile

2-{5'[spiro[cyclohexane-1,3'-[3H]indol]-(1'H)-one-2'(O-benzyloxime)]}-1H-pyrrole-1-carboxylic acid, *tert*-butyl ester. A solution of 5'-bromospiro{cyclohexane-1,3'-[3H]indol}-2' (1'H)-one 2'(O-benzyloxime) (7.4 g, 19.17 mmol) and tetrakis (triphenylphosphine)palladium (0) (2.5 g, 2.00 mmol) in DME (100 ml) was stirred under nitrogen for 15 minutes. To the solution was added 1-*tert*-butoxycarbonylpyrrole boronic acid (5.5 g, 26 mmol) and 1M sodium carbonate (50 ml). The mixture was heated to 80° C for 6 hours and allowed to cool. The reaction mixture was poured into water and extracted with ethyl acetate (3 x 100 ml). The organic layers were combined and dried over magnesium sulfate. The solution was filtered, concentrated *in vacuo*, and the residue was purified by flash chromatography on silica gel (4.5:1 Hexane/ ethylacetate) to give the product (7.7 g, 88 %) as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 1.28 (s, 9H), 1.55 - 1.66 (m, 8H), 1.83 - 1.98 (m, 2H), 4.99 (s, 2H), 6.12 - 6.14 (m, 1H), 6.22 (t, 1H, *J* = 3.26 Hz), 6.76 (d, 1H, *J* = 7.9 Hz), 7.02 (dd, 1H, *J* = 7.98, 1.4 Hz), 7.19 (s, 1H), 7.27 - 7.31 (m, 2H), 7.35 (t, 1H, *J* = 6.8 Hz), 7.43 (d, 1H, *J* = 8 Hz), 9.55 (s, 1H).

5'-(1-*tert*-Butoxycarbonyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one-2'-(O-benzyloxime)-1'-carboxylic acid, *tert*-butyl ester. To a solution of 2-{5'[spiro[cyclohexane-1,3'-[3H]indol]-(1'H)-one-2'(O-benzyloxime)]}-1H-pyrrole-1-carboxylic acid, *tert*-butyl ester (7.7 g, 16.38 mmol) in THF (anhydrous, 100 mL) was added sodium hydride (0.665 g, 17 mmol) after hydrogen evolution ceased di-*tert*-butyldicarbonate (10.9 g, 50 mmol) and DMAP (0.20 g) was added and the reaction stirred at 65 °C for 18 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layers were combined, and dried over magnesium sulfate. The solution was filtered, concentrated *in vacuo*, to give the product (9.0 g, 15.76 mmol) which was taken directly to the next step.

To a solution of 5'-(1-*tert*-Butoxycarbonyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one-2'-(O-benzyloxime)-1'-carboxylic acid, *tert*-butyl ester

(9.0 g, 15.76 mmol) in THF (anhydrous, 75 mL) at -78°C was added chlorosulfonyl isocyanate (1.55 mL, 17.54 mmol). After 90 minutes, DMF (21 mL, 275 mmol) was added and the reaction was allowed to warm to room temperature. The reaction was poured into water (200 mL) and extracted with ethyl acetate (2 X 100 mL). The organic layers combined and dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography on silica gel (10% ethyl acetate/ Hexane) gave 5'-(5-cyano-1-tert-butoxycarbonyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester (7.6 g, 82%) as a white powder. ^1H NMR (DMSO-*d*₆, 300 MHz) δ 1.30 (s, 9H), 1.38 (s, 9H), 1.58-1.83 (m, 8H), 1.72-1.73 (m, 2H), 5.0 (s, 2H), 6.44-6.45 (d, 1H, $J = 3.76$), 7.25-1.46 (m, 10H).

5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester. To a solution of 5'-(5-cyano-1-tert-butoxycarbonyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester (7.6 g, 3.25 g, 48 mmol) in THF (anhydrous, 30 mL) was added a solution of sodium ethoxide in ethanol (120 mL). The reaction mixture was heated to 80°C and stirred overnight. The mixture was cooled to room temperature and concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed with water, brine, and dried over magnesium sulfate. The solvent was evaporated *in vacuo* to afford the product (6.1 g, 95%). ^1H NMR (DMSO, 500 MHz) δ 1.38 (s, 9H), 1.63-1.74 (m, 8H), 1.88-1.97 (m, 2H), 5.08 (s, 2H) 6.69-6.7 (d, 1H, $J = .8\text{Hz}$), 6.98-6.99 (d, 1H, $J = .7\text{Hz}$), 7.29-7.37 (m, 1H), 7.35 (m, 2H), 7.42 (m, 3H), 7.63 (dd, 1H, $J = 1.8, .3\text{Hz}$), 7.76 (d, 1H, $J = .4\text{Hz}$).

5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester. To 5'-(5-cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(1'H)one-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester (6.1 g, 12.29 mmol) in DMF (75 mL) was added potassium carbonate (6.5 g, 47 mmol) and MeI (1 mL, 15.4 mmol) and the reaction mixture was stirred at room temperature for 2.5 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine and

the solvent was concentrated *in vacuo*. To give the desired product (6.1 g, 12.29 mmol) which was carried on to the next step without further purification. ¹H NMR (DMSO, 300 MHz) δ 1.38 (s, 9H), 1.62-1.98 (m, 10H), 3.71 (s, 3H), 5.08 (s, 2H), 6.34 (d, 1H, *J* = 4.1), 7.03 (d, 1H, *J* = 3.99), 7.30-7.53 (m, 8H).

5 5-{5'-Spiro[cyclohexane-1,3'-[3H]indol]-(1'H)-one-2'-(O-benzyloxime)}}-1H-pyrrole-1-methyl-2-carbonitrile. 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester (6.1 g, 12.29 mmol) was dissolved in dioxane (5 mL) and 4 M HCl in dioxane (10 mL) was added and the reaction heated to 45 °C for 3.5 hours. The
10 mixture was carefully neutralized with sodium bicarbonate (sat.). The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine and dried over magnesium sulfate. The solvent was evaporated *in vacuo*. Purification by column chromatography on silica gel (5% ethyl acetate/hexane) gave the product (4.36 g, 94%). ¹H NMR (DMSO-d₆, 300MHz) δ
15 1.57-1.7 (m, 8H), 1.9-2.05 (m, 2H), 3.68 (s, 3H), 5.00 (s, 2H), 6.25 (d, 1H, *J* = 3.92), 6.85 (d, 1H, *J* = 8.03), 7.00 (d, 1H, *J* = 4.08), 7.2-7.44 (m, 7H), 9.7 (s, 1H)

To 5-{5'-spiro[cyclohexane-1,3'-[3H]indol]-(1'H)-one-2'-(O-benzyloxime)}}-1H-pyrrole-1-methyl-2-carbonitrile (4.36g, 10.6 mmol) in methylene chloride (50 mL) was added 1M boron tribromide (35 mL, in methylene chloride) at -78° C. The
20 reaction mixture was allowed to warm to room temperature. After 4 hours, the reaction mixture was quenched with saturated sodium bicarbonate (100 mL). The organic layer was collected and the aqueous layer was extracted with ethyl acetate (2X 100 mL), organic layers combined, washed with brine, dried over magnesium sulfate, and the solvent evaporated *in vacuo*. The residue was purified by flashed
25 chromatography on silica gel (7:3 hexane/ethylacetate) to give the title compound (1.35g, 40%) as a white solid. ¹H NMR (DMSO-d₆, 300 MHz) δ 1.58 - 1.71 (m, 8H), 1.99 - 2.00 (m, 2H), 3.69 (s, 3H) 6.24 (d, 1H, *J* = 4.07 Hz), 6.8 (d, 1H, *J* = 8.05 Hz), 6.99 (d, 1H, *J* = 4.01 Hz), 7.20 (dd, 1H, *J* = 8.04, 1.57 Hz), 7.36 (d, 1H, *J* = 1.12 Hz), 9.48 (s, 1H), 9.62 (s, 1H).

Example 57**5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile**

5 5-(spiro[cyclohexane-1,3'-[3]indole]-2'(1H)-(O-benzyloxime))-1H-pyrrole-2-carbonitrile. Prepared from 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1H)-one-2'-(O-benzyloxime)-1'-carboxylic acid, tert-butyl ester (0.395 g, 0.796 mmol) dissolved in 2mL of THF and 4M HCl Dioxane/water (10 mL)
10 following the procedure used to prepare 5-{5'-spiro[cyclohexane-1,3'-[3H]indol]- (1'H)-one-2'-(O-benzyloxime)}}-1H-pyrrole-1-methyl-2-carbonitrile the desired product was obtained (0.220 g, 0.745 mmol, 95%). ¹ H NMR (DMSO-d₆, 500 MHz) δ 1.44 - 1.50 (m, 1H), 1.61 - 1.70 (m, 7H), 1.94 - 1.99 (m, 2H), 5.0 (s, 2H), 6.55 (d, 1H, J = 4 Hz), 6.79 (d, 1H, J = 8.0 Hz), 6.95 (d, 1H, J = 4 Hz), 7.27 - 7.31 (m, 1H),
15 7.34 - 7.37 (m, 2H), 7.42 - 7.43 (m, 2H), 7.47 (dd, 1H, J = 8.0, 1.4 Hz), 7.65 (d, 1H, J = 1.5 Hz), 9.65 (s, 1H), 12.4 (s, 1H).

 The title compound was prepared from 5-(spiro[cyclohexane-1,3'-[3]indole]- 2'(1H)-(O-benzyloxime))-1H-pyrrole-2-carbonitrile (0.325 g, 0.82mmol) and 1M
20 Boron tribromide (6mL in methylene chloride), following the procedure for 5-(spiro[cyclohexane-1,3'-[3H]indole]-2'-(hydroxyimino)-5'-yl)- 1H-pyrrole-1-methyl-2-carbonitrile, to obtain the product as an off white solid (0.110 g, 0.326 mmol, 44%).
¹ H NMR (DMSO-d₆, 500 MHz) δ 1.46 - 1.5 (m, 1H), 1.62 - 1.71 (m, 7H), 1.95 - 2.05 (m, 2H), 6.55 (d, 1H, J = 4.0 Hz), 6.75 (d, 1H, J = 8.0 Hz), 6.94 (d, 1H, J = 3.47 Hz), 7.45 (dd, 1H, J = 8.1, 1.73 Hz), 7.63 (d, 1H, J = 1.73), 9.42 (s, 1H), 9.59 (s, 1H),
25 12.39 (s, 1H).

Example 58**4-(Spiro[cyclohexane-1,3'-[3H]indole]-2'(acetoxyimino)-5'-yl)-2-thiophenecarbonitrile**

5

To a solution of 4-(Spiro[cyclohexane-1,3'-[3H]indole]-2'(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile (2.21 g, 6.83 mmol) and acetic anhydride (1ml) in dichloromethane-pyridine (30 ml, 9:1) was added 4-dimethylaminopyridine (250 mg) at room temperature. After 3h., the mixture was diluted with dichloromethane, washed with water, dil. Hydrochloric acid, water, dried (MgSO₄), and evaporated. The residue was purified by column chromatography (EtOAc: hexane, gradient elution) to afford the title compound (0.84 g, 2.29 mmol, 33%) as a white solid: MS (ESI (+ ve)) *m/z* 366 [M+H]⁺.

15

Example 59**3-Fluoro-N'-hydroxy-5-[2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl]benzenecarboximidamide**

5'-(3-Cyano-5-fluorophenyl)-2-(methylthio)spiro[cyclohexane-1,3'-[3H]indole]. Prepared from 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile (0.451 g, 1.34 mmol) according to the procedure described in Example 43 to afford the desired product (0.316 g, 0.90 mmol, 67%): ¹H NMR (DMSO, 300 MHz) δ 7.74 (d, 1H, *J* = 1.7 Hz), 7.68 (t, 1H, *J* = 1.4 Hz), 7.58 (d, 1H, *J* = 8.0 Hz), 7.54 (t, 1H, *J* = 2.3 Hz), 7.50 (dd, 1H, *J* = 8.0 and 1.9 Hz), 7.33 – 7.29 (m, 1H), 2.67 (s, 3H), 2.04 – 1.78 (m, 7H) and 1.58 – 1.50 (m, 3H); MS (ESI(+ ve)) *m/z* 351 (M+H)⁺.

25

To a solution of the last cited product (0.30 g, 0.88 mmol) in DMSO (10 ml) was added hydroxylamine (50% aqueous solution, 1 ml), and the reaction was heated to 120 °C. After 1h., the mixture was cooled, partitioned between saturated aqueous ammonium chloride and ethylacetate. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column

30

chromatography (SiO₂, 5% MeOH in dichloromethane) to afford the title compound (0.079 g, 0.23 mmol, 26%) as a white foam: ¹H NMR (DMSO, 300 MHz) δ 9.79 (s, 1H), 9.61 (s, 1H), 9.42 (s, 1H), 7.73 (s, 1H), 7.61 (d, 1H, *J* = 1.3 Hz), 7.46 (dd, 1H, *J* = 8.3 and 1.5 Hz), 7.34 (d, 1H, *J* = 10 Hz), 6.81 (d, 1H, *J* = 8.0 Hz), 6.01 (s, 2H), 2.11 – 2.02 (m, 2H) and 1.81 – 1.56 (m, 8H); MS (ESI(+ ve)) *m/z* 369 (M+H)⁺.

Example 60

N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indole]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide

10 4-Methyl-5-(spiro[cyclohexane-1,3'-[3H]indol] 2'-(methylthio)-5'-yl)-2-thiophenecarbonitrile. To potassium *tert*-butoxide (0.32g, 2.6 mmol) in THF was added 5-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-methyl-2-thiophenecarbonitrile (0.84 g, 2.5 mmol). After 15 minutes, methyl iodide (0.50 g, 3.48 mmol) was added. After 3 hours, reaction was poured into ammonium chloride
15 (sat.) and extracted with ethylacetate. The organic layers were combined and dried over magnesium sulfate. The solution was filtered, concentrated *in vacuo*, and the residue was purified by flash chromatography on silica gel (4:1 Hexane/ethyl acetate) to give the desired product (0.530 g, 85%). (DMSO, 300 MHz) δ 1.48 (m, 3H), 1.70 (m, 2H), 1.81 (m, 5H), 2.32 (s, 3H), 2.62 (s, 3H), 7.48 (dd, 1H, *J* = 7.87 Hz, 1.46 Hz),
20 7.5 (d, 1H, *J* = 8.05 Hz), 7.77 (d, 1H, *J* = 1.46 Hz), 7.88 (s, 1H).

To 4-methyl-5-(spiro[cyclohexane-1,3'[3H] indol]2'-(methylthio)-5'-yl)-2-thiophenecarbonitrile (.450 g, 1.3 mmol) in DMSO (1 mL) was added hydroxylamine hydrochloride (2 mL, 50% sol. in water) and heated to 100° C for 2.5 hours. Water added until solution became slightly turbid, allowed the mixture to cool to room
25 temperature. The white solid was filtered, collected and dissolved in ethyl acetate and dried over magnesium sulfate. The solution was filtered, concentrated *in vacuo*, giving (0.320 g, 69%). (DMSO-d₆, 500 MHz) δ 1.4-1.74 (m, 8H), 1.94-2.49 (m, 2H), 2.54 (s, 3H), 5.8 (s, 1H), 6.79 (d, 1H, *J* = 8.0), 7.16 (dd, 1H, *J* = 8.12, 1.83), 7.39 (m, 2H), 9.42 (s, 1H), 9.56 (s, 1H), 9.58 (s, 1H).

Example 61**N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indole]-2'-9hydroxyimino)-5'-yl-2-thiophenecarboximidamide**

4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(methylthio)-5'-yl]-2-thiophenecarbonitrile (0.077 g, .237 mmol) was reacted with 50% solution of hydroxylamine (1 mL) following the procedure for Example 60 to afford the title compound (0.016g, 0.044 mmol, 20 %). MS (ESI, (+VE)) m/z 357 [M+H]⁺.

Example 62**N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboxidamide**

The title compound was prepared from 5-(spiro[cyclohexane-1,3'-[3H]indol]2'-(methylthio)-5'-yl)-2-thiophenecarbonitrile (0.500 g, 1.5 mmol) and a 50% solution of hydroxylamine (2 mL, excess) following the procedure for Example 60, to afford the product (0.200 g, 0.56 mmol, 56%). ¹H NMR (DMSO-d₆, 500 MHz) δ 1.45-1.75 (m, 8H), 1.97-2.06 (m, 2H), 5.89 (s, 1H), 6.74 (d, 1H, J = 8 Hz), 7.3 (d, 1H, J = 3.9 Hz), 7.34 (dd, 1H, J = 8.06, 1.46 Hz), 7.4 (d, 1H, J = 8.0 Hz), 7.5 (d, 1H, 1.95 Hz), 9.44 (s, 1H), 9.58 (s, 1H), 9.6 (s, 1H).

Example 63**5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide**

5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-amine. To a turbid solution of 5'-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3-[3H]indol]-2-amine (0.500 g; 1.53 mmol) in 25 mL of ethanol was added hydrazine hydrate (0.600 mL; 12.24 mmol). The solution was warmed to 55°C, where Raney-nickel (50% in water) was added to the reaction to keep a constant evolution of gas. After 45 minutes, the hot reaction mixture was filtered through a Celite plug and rinsed with a copious amount of hot methanol. The filtrate was concentrated in vacuo to give 0.890 g of an opaque solid. The product was purified by flash silica gel chromatography; (eluant, 2% to 8 % methanol-methylene chloride with 0.1 % ammonium hydroxide)

to afford 0.310 g (65%) of the desired product as a white solid. Mp. 118-120° C. ¹H NMR δ (300 MHz, DMSO-d₆) 1.31-1.46 (m, 2H), 1.70-1.93 (m, 8H), 7.0 (d, 1H), 7.1 (br, 2H, 2NH), 7.31-7.34 (dt, 1H, J= 8 Hz), 7.41-7.46 (t, 2H), 7.55-7.58 (d, 1H), 7.62 (s, 1H), 7.72 (s, 1H); MS (ECI(+ve)) m/z 311 (M+H)⁺.

- 5 1-tert-butoxycarbonyl-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-amine. To a solution of 5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-amine (0.310 g; 0.96 mmol) in dry methylene chloride at 0° C was added Di-tert-butyl dicarbonate (0.252 g; 1.15 mmol) and 4-dimethylaminopyridine (0.117 g; 0.96 mmol). The solution was allowed to warm to room temperature and stir 24 h. The
10 reaction solution was diluted with water (50 mL) and the layers were separated. The organic layer was dried over Na₂SO₄, filtered and concentrated *in vacuo* to give 0.355 g of a yellow oil. The product was purified by flash silica gel chromatography; (eluant, 1% to 3 % methanol- methylene chloride) to afford the desired product (0.081 g, 20%) as a white solid. ¹H NMR δ (300 MHz, DMSO-d₆) 1.58 (m, 2H), 1.63 (s, 9H,
15 Boc), 1.77-1.79 (m, 8H), 7.42-7.48 (m, 2H), 7.64-7.68 (m, 3H), 7.70-7.80 (m, 2H), 9.72 (s, 1H, NH). MS (ECI(+ve)) m/z 411 (M+H)⁺.


- 1'-tert-Butoxycarbonyl-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-amine (0.120 g; 0.29 mmol) in 2.0 mL of dry DMF was added to a solution of 4- dimethylaminopyridine (0.089 g; 0.73 mmol) and cyanogen bromide
20 (0.077 g; 0.73 mmol) in 4.0 mL of dry DMF at 0° C. The yellow solution was heated to 40° C for 16 h. Work-up included pouring the reaction solution into 0.1 N NaHCO₃ (50 mL) and extracting with ethyl acetate (3 x 50 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo* to give 0.091 g of a yellow residue. The product was purified by flash silica gel chromatography;
25 (stepwise gradient of 5:1 to 3:1 hexane : ethyl acetate) to afford 0.031 g (32%) of the product as a bright yellow solid. Mp. 225°C (dec.). ¹H NMR δ (500 MHz, DMSO-d₆) 1.46-1.73 (m, 8H), 1.89-1.90 (m, 2H), 7.13-7.16 (d, 1H), 7.38-7.41 (dt, 1H, J= 8Hz), 7.45-7.50 (m, 1H), 7.60-7.63 (dd, 2H, J= 6.4 Hz), 7.71 (s, 1H), 7.85 (s, 1H), 12.1 (s, 1H, NH); MS (ECI(-ve)) m/z 336 (M-H)⁻.

Other desirable compounds, which can be made according to the methods described herein, include 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide, 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide, 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide, 5'-(5-Cyano-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-cyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol-5'-yl])-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol-5'-yl])-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol-5'-yl])-1-methyl-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol-5'-yl])-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol-5'-yl])-4-methyl-thiophene-2-carbonitrile, 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol-5'-yl])-thiophene-2-carbonitrile.

All publications cited in this specification are incorporated herein by reference herein. While the invention has been described with reference to a particularly preferred embodiment, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

What is Claimed:

A compound of the formula: 1 or 2:



wherein:

R_1 and R_2 are chosen independently from the group of H, alkyl, substituted alkyl; OH; O(alkyl); O(substituted alkyl); OAc; aryl; optionally substituted aryl; heteroaryl; optionally substituted heteroaryl; alkylaryl; alkylheteroaryl; 1-propynyl; or 3-propynyl:

or R_1 and R_2 are joined to form a ring comprising one of the following:

$-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$; $-\text{CH}_2\text{CH}_2\text{CMe}_2\text{CH}_2\text{CH}_2-$; $-\text{O}(\text{CH}_2)_m\text{CH}_2-$; $\text{O}(\text{CH}_2)_p\text{O}-$;
 $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$; or $-\text{CH}_2\text{CH}_2\text{N}(\text{H or alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R_1 and R_2 together comprise a double bond to one of the following:

CMe_2 ; C(cycloalkyl), O, C(cyloether).

R_3 is selected from H, OH, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_6 alkenyl, alkynyl or substituted alkynyl, or COR^A ;


R^A is selected from H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, or substituted C_1 to C_3 aminoalkyl;

- 111 -

R_4 is selected from H, halogen, CN, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_1 to C_6 alkoxy, substituted C_1 to C_6 alkoxy, C_1 to C_6 aminoalkyl, or substituted C_1 to C_6 aminoalkyl;

R^5 is selected from the groups a), b) or c):

a) R^5 is a trisubstituted benzene ring containing the substituents X, Y and Z as shown below:



X is selected from halogen, OH, CN, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 thioalkyl, substituted C_1 to C_3 thioalkyl, $S(O)alkyl$, $S(O)_2alkyl$, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 or 6 membered heterocyclic ring containing 1 to 3 heteroatoms, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CON(alkyl)_2$, $CSN(alkyl)_2$, COR^B , $OCOR^B$, $NR^C COR^B$;

R^B is selected from H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, or substituted C_1 to C_3 aminoalkyl;

R^C is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

Y and Z are independently selected from H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_3 alkyl, or C_1 to C_3 thioalkyl;

or

b) R^5 is a five or six membered heterocyclic ring with 1, 2, or 3 heteroatoms selected from O, S, SO, SO_2 or NR^D and containing one or two independent substituents from the group of H, halogen, CN, NO_2 and C_1 to C_3 alkyl, C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, COR^D , or $NR^E COR^D$;

R^D is H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, or substituted C_1 to C_3 aminoalkyl;

- 112 -

R^E is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

R^6 is H, or C_1 to C_3 alkyl; or

c) R^5 is an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, the moiety being optionally substituted by from 1 to 3 substituents selected from halogen, lower alkyl, CN, NO_2 , lower alkoxy, or CF_3 ;


Q^1 is S, NR_7 , CR_8R_9 ;

R_7 is selected from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, acyl, substituted acyl, aroyl, substituted aroyl, SO_2CF_3 , OR^{11} or $NR^{11}R^{12}$;


R_8 and R_9 are independent substituents selected from the group of H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN, or CO_2R_{10} ,

R_{10} is C_1 to C_3 alkyl; or

CR_8R_9 comprises a six membered ring as shown by the structure below




Q^2 is selected from the moieties:




R^{11} , R^{12} and R^{13} are independently selected from H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, aroyl or substituted aroyl or sulfonyl; or a pharmaceutically acceptable salt thereof.

2. A compound of Claim 1 having the formula:



wherein R₅ is a disubstituted benzene ring of the formula:



X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy; and


Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; and

Q¹ is S, NR₇, CR₈R₉;


or a pharmaceutically acceptable salt thereof.

3. A compound of Claim 1 having the formula:

- 114 -



wherein R_5 is a five membered ring with the structure shown below



wherein:

U is O, S, or NR_6 ;

R_6 is H, or C_1 to C_3 alkyl, or C_1 to C_4 CO_2 alkyl;


X' is selected from halogen, CN, NO_2 , $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl_2$, $CSNalkyl_2$, C_1 to C_3 alkyl, or C_1 to C_3 alkoxy;

Y' is from the group of H, F or C_1 to C_4 alkyl; and

Q^1 is S, NR_7 , CR_8R_9 ;


or a pharmaceutically acceptable salt thereof.

4. A compound of Claim 1 having the formula:



wherein R_5 is a six membered ring with the structure shown

- 115 -



wherein X^1 is N or CX^2 ,

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl_2$, $CSNalkyl_2$ or NO_2 ;


Q^1 is S, NR_7 , CR_8R_9 ;

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN , CO_2R_{10} .

R_{10} is C_1 to C_3 alkyl;

CR_8R_9 are within a six membered ring as shown by the structure below



or pharmaceutically acceptable salt thereof

6. A compound of Claim 1 which is 5'-(3-Chlorophenyl)spiro [cyclohexane-1,3'-[3H]indole]-2'(1'H)-thione or a pharmaceutically acceptable salt thereof.

- 116 -


7. A compound of Claim 1 which is 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol-5'-yl) benzonitrile or a pharmaceutically acceptable salt thereof.

8. A compound of Claim 1 which is 4-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile or a pharmaceutically acceptable salt thereof.


9. A compound of Claim 1 which is 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile or a pharmaceutically acceptable salt thereof.

10. A compound of Claim 1 which is 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophene thioamide or a pharmaceutically acceptable salt thereof.

11. A compound of the formula:



each wherein R₅ is a disubstituted benzene ring containing the substituents X and Y:




- 117 -


X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

12. A compound according to claim 11, having the formulae:




are those wherein R₅ is a six membered ring with the structure:



wherein X¹ is N or CX²,

X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂ or NO₂;

Q² is selected from the moieties:




R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN CO_2R_{10} ,



R_{10} is C_1 to C_3 alkyl;

CR_8R_9 are within a six membered ring as shown by the structure below



or pharmaceutically acceptable salt thereof.


13. A compound of the formula:



wherein R_{14} is chosen from the group H, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, substituted sulfonyl.

wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below


- 119 -



X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNHOH, CNH₂NOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

14. A compound according to claim 13, wherein R₅ is a five membered ring with the structure shown below



wherein:

U is O, S, or NR₆;

R₆ is H, or C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;


X' is selected from halogen, CN, NO₂, CONH₂, CNHNHOH, CNH₂NOH, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkyl, or C₁ to C₃ alkoxy;

Y' is from the group of H, F or C₁ to C₄ alkyl; or a pharmaceutically acceptable salt thereof.

15. The compound according to claim 13, wherein R₅ is a thiophene or furan ring substituted by X' and Y'.

- 120 -

16. The compound according to claim 13, wherein R_5 is a six membered ring with the structure:



wherein X^1 is N or CX^2 ,

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl_2$, $CSNalkyl_2$ or NO_2 ;


R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN , CO_2R_{10} ,

R_{10} is C_1 to C_3 alkyl;


or pharmaceutically acceptable salt thereof.

17. A compound of the formula:



wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below


- 121 -



X is selected from halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, CNHNOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C₁ to C₃ thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, or C₁ to C₃ thioalkyl; or a pharmaceutically acceptable salt thereof.

18. The compound according to claim 17, wherein R₅ is a five membered ring with the structure shown below



wherein:

U is O, S, or NR₆;

R₆ is H, or C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkyl, or C₁ to C₃ alkoxy;


Y' is from the group of H, F or C₁ to C₄ alkyl;

or a pharmaceutically acceptable salt thereof.

19. The compound according to claim 17, wherein R₅ is a thiophene or furan ring substituted by X' and Y', as described above.

- 122 -

20. The compound according to claim 17, wherein R_5 is a six membered ring with the structure:



wherein X^1 is N or CX^2 ,

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl_2$, $CSNalkyl_2$ or NO_2 ;

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;


R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN , CO_2R_{10} ,

R_{10} is C_1 to C_3 alkyl;

or pharmaceutically acceptable salt thereof.


- 123 -

21. A compound of the formula:



wherein R_{15} is selected from the group H, Me, CO_2R , acyl, substituted acyl, aroyl, substituted aroyl, alkyl, substituted alkyl, CN;

wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below




X is selected from halogen, CN, CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CONalkyl_2 , CSNalkyl_2 , CNHNOH , C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C_1 to C_3 thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, or C_1 to C_3 thioalkyl; or a pharmaceutically acceptable salt thereof.

22. The compound according to claim 21, R_5 is a five membered ring with the structure shown below

- 124 -



wherein:

U is O, S, or NR₆;

R₆ is H, or C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;


X' is selected from halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂, C₁ to C₃ alkyl, or C₁ to C₃ alkoxy;

Y' is from the group of H, F or C₁ to C₄ alkyl;

or a pharmaceutically acceptable salt thereof.

23. The compound according to claim 21, wherein R₅ is a thiophene or furan ring substituted by X' and Y'.

24. The compound according to claim 21, wherein R₅ is a six membered ring with the structure:



wherein X' is N or CX²,

X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CONalkyl₂, CSNalkyl₂ or NO₂;


R₇ is from the group including CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO₂CF₃;

R₈ and R₉ are independent substituents from the group including H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO₂, CN CO₂R₁₀,


- 125 -

R_{10} is C_1 to C_3 alkyl;
or pharmaceutically acceptable salt thereof.

25. A compound of the formula:



wherein R_5 is a disubstituted benzene ring containing the substituents X and Y as shown below




X is selected from halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CONalkyl$, $CSNalkyl$, $CNHNOH$, C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring containing 1 to 3 heteroatoms, or C_1 to C_3 thioalkoxy;

Y is a substituent on the 4' or 5' position from the group including H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, or C_1 to C_3 thioalkyl;
or a pharmaceutically acceptable salt thereof.

26. The compound according to claim 25, wherein R_5 is a five membered ring with the structure shown below

- 126 -



wherein:

U is O, S, or NR_6 ;

R_6 is H, or C_1 to C_3 alkyl, or C_1 to C_4 CO_2 alkyl;


X' is selected from halogen, CN, NO_2 , CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CONalkyl_2 , CSNalkyl_2 , C_1 to C_3 alkyl, or C_1 to C_3 alkoxy;

Y' is from the group of H, F or C_1 to C_4 alkyl;

or a pharmaceutically acceptable salt thereof.

27. The compound according to claim 26, wherein R_5 is a thiophene or furan ring substituted by X' and Y' .

28. The compound according to claim 26, wherein R_5 is a six membered ring with the structure:



wherein X' is N or CX^2 ,

X^2 is halogen, CN, CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CONalkyl_2 , CSNalkyl_2 or NO_2 ;

R_7 is from the group including CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, or SO_2CF_3 ;

R_8 and R_9 are independent substituents from the group including H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, NO_2 , CN , CO_2R_{10} ,

R_{10} is C_1 to C_3 alkyl;
or pharmaceutically acceptable salt thereof.

29. A pharmaceutical composition comprising a compound according to any of claims 1 to 28, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

30. A method of inducing contraception in a mammal, the method comprising administering to a mammal in need thereof a pharmaceutically effective amount of a compound of any of claims 1 to 28, or a pharmaceutically acceptable salt thereof.

31. A method of treatment of dysfunctional bleeding in a mammal, the method comprising administering to a mammal in need thereof a pharmaceutically effective amount of a compound of any of claims 1 to 28, or a pharmaceutically acceptable salt thereof.

32. A method of treatment of carcinomas and adenocarcinomas of the endometrium, ovary, breast, colon, or prostate in a mammal, the method comprising administering to a mammal in need thereof a pharmaceutically effective amount of a compound of any of claim 1 to 28, or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

Intern 1st Application No

PCT/US 00/11630

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/08 C07D209/30 C07D209/40 A61K31/404 A61P5/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EP0-Internal, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 27059 A (MEIJI SEIKA KAISHA, LTD) 25 June 1998 (1998-06-25) claim 1	1,29
P,A	& EP 0 947 507 A 6 October 1999 (1999-10-06) page 59 -page 70	
A	WO 97 13767 A (CHEMISCH PHARMAZEUTISCHE FORSCHUNGSGESELLSCHAFT) 17 April 1997 (1997-04-17) cited in the application page 1	1,29
A	WO 91 06545 A (BOEHRINGER MANNHEIM GMBH) 16 May 1991 (1991-05-16) cited in the application page 1 -page 5, paragraph 1	1,29
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 September 2000

Date of mailing of the international search report

15/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/11630

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 29272 A (ASTRA AKTIEBOLAG) 22 December 1994 (1994-12-22) page 1 -page 3 -----	1,29
A	US 4 721 721 A (DONALD E. KUHLA ET AL.) 26 January 1988 (1988-01-26) the whole document -----	1,29
A	EP 0 208 510 A (PFIZER INC.) 14 January 1987 (1987-01-14) page 1 -----	1,29
A	ALDO ANDREANI ET AL.: "Potential antitumor agents XVII" ACTA PHARM. NORD., vol. 2, no. 6, 1990, pages 407-414, XP000925899 cited in the application the whole document -----	1,29

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/11630

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9827059 A	25-06-1998	AU 7738098 A BR 9714145 A EP 0947507 A US 6077840 A	15-07-1998 29-02-2000 06-10-1999 20-06-2000
WO 9713767 A	17-04-1997	AU 7284096 A	30-04-1997
WO 9106545 A	16-05-1991	DE 3935514 A AT 131821 T AU 7042191 A DE 59009988 D EP 0497852 A	02-05-1991 15-01-1996 31-05-1991 01-02-1996 12-08-1992
WO 9429272 A	22-12-1994	AU 7010894 A CA 2164119 A EP 0703901 A FI 956074 A JP 8511515 T NO 955074 A	03-01-1995 22-12-1994 03-04-1996 18-12-1995 03-12-1996 07-02-1996
US 4721721 A	26-01-1988	AU 5238886 A EP 0207140 A WO 8603749 A US 4818755 A	22-07-1986 07-01-1987 03-07-1986 04-04-1989
EP 208510 A	14-01-1987	US 4678802 A AT 67185 T AU 566065 B AU 5984786 A CA 1286663 A CN 86105309 A DD 251971 A DE 3681358 A DK 324286 A ES 2001855 A ES 2009227 A FI 862882 A GR 861760 A HU 41386 A IE 58566 B IL 79356 A JP 1883341 C JP 6010193 B JP 62026269 A KR 8802272 B NO 862750 A NZ 216788 A PH 22641 A PT 82931 A, B US 4725616 A US 4791131 A YU 114487 A YU 120386 A ZA 8605019 A US 4730004 A	07-07-1987 15-09-1991 08-10-1987 09-04-1987 23-07-1991 14-01-1987 02-12-1987 17-10-1991 10-01-1987 01-07-1988 16-09-1989 10-01-1987 10-11-1986 28-04-1987 06-10-1993 19-03-1990 10-11-1994 09-02-1994 04-02-1987 21-10-1988 12-01-1987 28-10-1988 28-10-1988 01-08-1986 16-02-1988 13-12-1988 29-02-1988 29-02-1988 24-02-1988 08-03-1988